

Administration of COVID-19 mRNA vaccine to individuals from 6 months of age

Patient group direction (PGD) template

Publication date: 24th September 2026

Expiry date: 30th April 2027

PGD Number: 2026/2980

Version 1.0



Translations



Easy read



BSL



Audio



Large print



Braille

Translations and other formats are available on request at:

 phs.otherformats@phs.scot

Public Health Scotland is Scotland's national agency for improving and protecting the health and wellbeing of Scotland's people.

© Public Health Scotland 2026



This publication is licensed for re-use under the [Open Government Licence v3.0](#).

For more information, visit
www.publichealthscotland.scot/ogl

www.publichealthscotland.scot

Most recent changes

Version	Date	Summary of changes
1.0	24 September 2026	New combined PGD template to support delivery of the Autumn 2026 COVID-19 vaccination programme to eligible individuals aged 6 months and over. This PGD reflects the change in antigenic content of the COVID-19 vaccine, from LP.8.1 (as utilised in Spring 2026 campaign) to XFG. The PGD also amalgamates the 2 previously separate PGDs for adults and children aged 6 months to 17 years, into a single legal framework.

Contents

Most recent changes	1
Authorisation	3
1. Clinical situation	4
2. Description of treatment	12
3. Adverse reactions	22
4. Characteristics of staff authorised under the PGD	27
5. Audit trail	30
6. Additional references	31
7. PGD for administration of COVID-19 mRNA vaccine to individuals from 6 months of age version 1.0 (valid from 24 September 2026 and expires 30 April 2027): authorisation	33
8. Version history	35

Authorisation

PGD for administration of COVID-19 mRNA vaccine to individuals from 6 months of age



This specimen Patient Group Direction (PGD) template has been produced by Public Health Scotland to assist NHS Boards. NHS Boards should ensure that the final PGD is considered and approved in line with local clinical governance arrangements for PGDs.

The registered health professionals who may administer COVID-19 mRNA vaccine can only use this PGD as named individuals. It is the responsibility of each professional to practice within the bounds of their own competence and in accordance with their own Code of Professional Conduct and to ensure familiarity with the manufacturer's product information/Summary of Product Characteristics (SmPC) for the vaccine administered in accordance with this PGD. NHS Board governance arrangements should indicate how records of staff authorised to operate this PGD will be maintained. Under PGD legislation there can be no delegation. Administration of the vaccine must be by the same practitioner who has assessed the patient under the PGD.

This PGD has been produced for NHS Greater Glasgow & Clyde by:

Professionals drawing up PGD/Authors	
*Name: Sarah Marshall  Signature: Date: 29/09/2026	Designation: Specialist Pharmacist Public Health E-mail address: sarah.marshall4@nhs.scot
Name: Tina McMichael  Signature: Date: 29/09/2026	Designation: Health Protection Nurse Specialist E-mail address: tina.mcmichael2@nhs.scot
Name: Iain Kennedy  Signature: Date: 29/09/2026	Designation: Consultant in Public Health E-mail address: iain.kennedy5@nhs.scot

Approved on behalf of NHS Greater Glasgow and Clyde by ADTC Patient Group
Direction Sub Committee representatives:

Chair	Dr Craig Harrow	
Director of Pharmacy representative	Elaine Paton, Senior Prescribing Adviser	
Director of Nursing Representative	Kevin McAuley, Chief Nurse North Sector	

Date approved: 29th September 2026

Valid from: 24th September 2026

Expiry date: 30th April 2027

1. Clinical situation

1.1. Indication

COVID-19 mRNA vaccines are indicated for active immunisation against COVID-19 disease caused by SARS-CoV-2 virus in those aged 6 months and over in accordance with Scottish Government COVID-19 immunisation programme and JCVI advice/recommendations as set out in **The Green Book COVID-19 Chapter** and subsequent correspondence/publications from **Scottish Government**.

1.2. Inclusion criteria

COVID-19 mRNA vaccine should be offered to those individuals aged 6 months and over, invited, or eligible in accordance with the recommendations in **The Green Book COVID-19 Chapter**, and/or in line with subsequent correspondence/publications from **Scottish Government**.

Individuals who have received a haematopoietic stem cell transplant or CAR-T therapy and who require revaccination, in accordance with the **Revaccination of patients following haematopoietic stem cell transplant or CAR-T treatment Guidance**.

Valid consent has been given to receive the vaccine.

1.3. Exclusion criteria

Individuals who:

- are aged under 6 months.
- have had a confirmed anaphylactic reaction to a previous dose of a mRNA COVID-19 vaccine.
- have had a confirmed anaphylactic reaction to any component of the vaccine or residual products from manufacture, these include polyethylene glycol (PEG). Practitioners must check the marketing authorisation holder's SmPC for details of vaccine components.
- have a history of immediate anaphylaxis to multiple, different drug classes, with the trigger unidentified (this may indicate PEG allergy) unless the advice from relevant specialist, local immunisation or health protection team is that vaccination should proceed.
- have a history of anaphylaxis to a vaccine, injected antibody preparation or a medicine likely to contain PEG (e.g. depot steroid injection, laxative) unless the advice from relevant specialist, local immunisation or health protection team is that vaccination should proceed.
- have a history of idiopathic (unexplained) anaphylaxis unless the advice from relevant specialist, local immunisation or health protection team is that vaccination should proceed.

- have evidence of current deterioration of COVID-19 symptoms: deferral of vaccination may be considered to avoid incorrect attribution of any change in the person's underlying condition to the vaccine.
- are suffering from acute severe febrile illness (the presence of a minor infection is not a contraindication for immunisation)
- are bone marrow and peripheral blood stem cell donors who have commenced Granulocyte-colony stimulating factor (GCSF): the vaccination (first or second dose) must be delayed at least until 72 hours after stem cell collection (both peripheral blood stem cell and bone marrow donation). This is precautionary advice to avoid vaccination when receiving GCSF and allow for post-donation recovery period
- have developed myocarditis or pericarditis following a previous dose of COVID-19 vaccination.

Under this PGD, some COVID-19 vaccines are restricted to use in particular age groups.

Practitioners must be familiar with and refer to the marketing authorisation holder's SmPC for the particular vaccine.

- Spikevax® XFG 0.1mg/1ml (50 micrograms/0.5ml dose) dispersion for injection COVID-19 mRNA vaccine is recommended for those aged 18 years and over.
- Comirnaty® XFG 30 micrograms/dose dispersion for injection in pre-filled syringe COVID-19 mRNA vaccine is recommended for those aged 12 to 17 years inclusive.
- Comirnaty® XFG 10 micrograms/dose dispersion for injection COVID-19 mRNA vaccine is recommended for those aged 6 months to 11 years inclusive.

1.4. Cautions/need for further advice/circumstances when further advice should be sought

The Green Book advises that there are very few individuals who cannot receive COVID-19 vaccines. Where there is doubt, rather than withholding vaccination, appropriate advice should be sought from the relevant specialist, or from the local immunisation or health protection team.

Individuals with a history of allergy

Those with a personal history of allergy should be managed in line with **The Green Book, COVID-19 Chapter**.

Where individuals have experienced a possible allergic reaction to a dose of COVID-19 vaccine, follow the guidance in the flowchart in **The Green Book COVID-19 Chapter** in relation to administration of subsequent doses.

The Green Book COVID-19 Chapter states individuals with non-allergic reactions (vasovagal episodes, non-urticarial skin reaction or non-specific symptoms) to the first dose of a COVID-19 vaccine can receive the second dose of vaccine in any vaccination setting. Observation for 15 minutes is recommended.

No specific management is required for individuals with a family history of allergies.

Individuals with thrombocytopenia

For individuals with a history of idiopathic thrombocytopenic purpura (ITP) receiving any COVID-19 vaccine, checking their platelet count should be considered 2 to 5 days post vaccination. Individuals who previously experienced ITP after the first dose of AstraZeneca® vaccine should be assessed by a haematologist and the risk benefit of further vaccination and with which COVID-19 vaccine should be considered on an individual basis.

Guillain-Barré syndrome (GBS)

Very rare reports have been received of GBS following COVID-19 vaccination. Healthcare professionals should be alert to the signs and symptoms of GBS to ensure correct diagnosis and to rule out other causes, in order to initiate adequate supportive care and treatment.

Individuals who have a history of GBS should be vaccinated as recommended for their age and underlying risk status. In those who are diagnosed with GBS after the first dose of vaccine, the balance of risk benefit is in favour of completing a full COVID-19 vaccination schedule. Where GBS occurs following either of the mRNA vaccines, further vaccination can proceed as normal, once recovered.

Capillary leak syndrome (applies to Spikevax® XFG Vaccine in this PGD)

Extremely rare reports of capillary leak syndrome have been reported after Spikevax® vaccines in individuals with a prior history of this condition. Individuals with a history of capillary leak syndrome, should be carefully counselled about the risks and benefits of vaccination and advice from a specialist should be sought.

Co-administration with other vaccines

Where individuals in an eligible cohort present having recently received another inactivated or live vaccine, COVID-19 vaccination should still be given. The same applies for other live and inactivated vaccines where COVID-19 vaccination has been received first or where a patient presents requiring 2 or more vaccines. It is generally better for vaccination to proceed to avoid any further delay in protection and to avoid the risk of the patient not returning for a later appointment. This includes, but is not limited, to vaccines commonly administered around the same time or in the same settings. Examples include LAIV, HPV, influenza, MenACWY and Td-IPV vaccines in the school age programmes, influenza, RSV and pertussis in pregnancy and influenza, pneumococcal, shingles and RSV vaccines in adults.

When administering at the same time as other vaccines, care should be taken to ensure that the appropriate route of injection is used for all the vaccinations. The vaccines should be given at separate sites, preferably in different limbs. If given in the same limb, they should be given at least 2.5cm apart. The site at which each vaccine was given should be noted in the individual's records.

Syncope

Syncope (fainting) can occur following, or even before, any vaccination especially in adolescents as a psychogenic response to the needle injection. This can be accompanied by several neurological signs such as transient visual disturbance, paraesthesia and tonic-clonic limb movements during recovery. It is important that procedures are in place to avoid injury from faints.

Pregnancy and breastfeeding

There is no known risk associated with giving inactivated, recombinant viral or bacterial vaccines or toxoids during pregnancy or whilst breastfeeding. Since inactivated vaccines cannot replicate, they cannot cause infection in either the mother or the foetus.

Vaccination should be offered for eligible pregnant women in accordance with recommendations in [The Green Book COVID-19 Chapter](#), following a discussion of the risks and benefits of vaccination with the woman.

There is no known risk associated with giving non-live vaccines whilst breastfeeding. JCVI advises that breastfeeding women may be offered vaccination with any suitable COVID-19 vaccine. Emerging safety data is reassuring: mRNA was not detected in the breast milk of recently vaccinated and protective antibodies have been detected in breast milk. The developmental and health benefits of breastfeeding should be considered along with the woman's clinical need for immunisation against COVID-19.

Individuals who have received COVID-19 vaccine overseas or as part of clinical trials

Eligible individuals who have been vaccinated abroad or in clinical trials should be offered COVID-19 vaccination in accordance with current recommendations. It is not necessary to know which vaccines an individual has previously received unless they indicate that they had an adverse event to a COVID-19 vaccine that would either contradict or be a precaution for a further dose. Specific advice is available from [UKHSA](#).

Individuals with a past history of COVID-19 infection

There are no safety concerns from vaccinating with a past history of COVID-19 infection, or with detectable COVID-19 antibody.

Vaccination of individuals who may be infected or asymptomatic or incubating COVID-19 infection is unlikely to have a detrimental effect on the illness although individuals with suspected COVID-19 infection should not attend vaccination sessions to avoid infecting others.

There is no need to defer immunisation in individuals after recovery from a recent episode with compatible symptoms, whether or not they are tested for COVID-19.

Having prolonged COVID-19 symptoms is not a contraindication to receiving COVID-19 vaccine but if the patient is seriously debilitated, still under active investigation, or has evidence of recent deterioration, deferral of vaccination may be considered to avoid incorrect attribution of any change in the person's underlying condition to the vaccine.

During care home outbreaks, vaccination of residents with confirmed COVID-19 may go ahead provided the residents are clinically stable and infection control procedures can be maintained.

1.5 Action if excluded

Specialist advice must be sought on the vaccine and circumstances under which it could be given. Immunisation using a patient specific direction may be indicated. The risk to the individual of not being immunised must be taken into account.

For individuals aged 12 years and over, where mRNA products are considered **not** clinically suitable, vaccination with a non mRNA product may be considered. Please refer to Administration of Bimervax LP.8.1 (Hipra COVID-19 Vaccine (recombinant, adjuvanted)) vaccine PGD.

For those aged less than 12 years where mRNA products are considered not clinically suitable, specialist advice should be sought for further assessment on alternative vaccine suitability.

Document the reason for exclusion and any action taken in accordance with local procedures.

Inform or refer to the clinician in charge.

Temporary exclusion

In case of postponement due to acute severe febrile illness or COVID-19 symptoms, advise when the individual can be vaccinated. Ensure another appointment is arranged or the individual knows how future vaccination may be accessed.

1.6. Action if patient declines

Advise the individual about the protective effects of the vaccine, the risks of infection and potential complications of disease.

Advise how future immunisation may be accessed if they subsequently decide to receive the vaccine.

Document advice given and decision reached.

Inform or refer to the clinician in charge.

2. Description of treatment

2.1. Name of medicine/form/strength

Spikevax® XFG 0.1mg/1ml dispersion for injection. Multidose vial that contains 5 doses of 0.5 ml. One dose (0.5 ml) contains 50 micrograms of mRNA-1273.262, a COVID-19 mRNA Vaccine (embedded in SM-102 lipidnanoparticles).

Comirnaty® XFG 30 micrograms/dose dispersion for injection in pre-filled syringe. One dose (0.3 mL) contains 30 micrograms of mRNA encoding XFG, a COVID-19 mRNA Vaccine (nucleoside modified, embedded in lipid nanoparticles).

Comirnaty® XFG 10 micrograms/dose dispersion for injection. Single dose vial. One dose (0.3 mL) contains 10 micrograms of mRNA encoding XFG, a COVID-19 mRNA Vaccine (nucleoside modified, embedded in lipid nanoparticles).

2.2. Route of administration

COVID-19 vaccines must be administered by intramuscular (IM) injection preferably into the deltoid area of the upper arm. Where administration into the deltoid is not possible the anterolateral thigh can be considered. For small infants the anterolateral thigh may be used.

Individuals with bleeding disorders may be vaccinated intramuscularly if, in the opinion of a doctor familiar with individual's bleeding risk, vaccines or similar small volume intramuscular injections can be administered with reasonable safety by this route. If the individual receives medication/ treatment to reduce bleeding, for example treatment for haemophilia, intramuscular vaccination can be scheduled shortly after such medication/treatment is administered. Individuals on stable anticoagulation

therapy, including individuals on warfarin who are up-to-date with their scheduled INR testing and whose latest INR is below the upper level of the therapeutic range, can receive intramuscular vaccination. A fine needle (23 or 25 gauge) should be used for the vaccination, followed by firm pressure applied to the site without rubbing for at least 2 minutes. The individual, parent or carer should be informed about the risk of haematoma from the injection.

Each vial contains at least the number of doses stated. It is normal for liquid to remain in the vial after withdrawing the final dose.

Care should be taken to ensure a full dose is administered. Where a full dose from a multi-dose vial cannot be extracted, the remaining volume should be discarded.

The vaccine should be visually inspected for particulate matter and discoloration prior to administration. In the event of any foreign particulate matter and/or variation of physical aspect being observed, do not administer the vaccine.

Vaccine unsuitable for use should be discarded in accordance with local procedures.

2.3. Dosage

Individuals aged over 18 years of age

Spikevax® XFG 0.1 mg/mL dispersion for injection COVID-19 mRNA Vaccine

Vaccine dose is 0.5ml (50 micrograms).

Individuals aged 12 years to 17 years

Comirnaty® XFG 30 micrograms/dose dispersion for injection COVID-19 mRNA

Vaccine dose is 0.3ml.

Infants and children aged 6 months to 11 years

Comirnaty® XFG 10 micrograms/dose dispersion for injection COVID-19 mRNA

Vaccine dose is 0.3ml.

2.4. Frequency

Individuals aged 5 years and over

A single dose regardless of prior COVID-19 vaccination status.

For individuals who have previously been vaccinated with a COVID-19 vaccine, the dose should be given at least 3 months (12 weeks) after the previous dose of COVID-19 vaccine (regardless of the vaccine given for the previous dose).

The only exception to the 3 months interval would be where individuals were about to receive or increase the intensity of an immunosuppressive treatment, and therefore a better response would be made if immunised prior to that treatment commencing. In this unusual scenario, the interval may be reduced to a minimum of 3 weeks on specialist clinical advice.

Individuals aged 6 months to 4 years

Primary immunisation

Comirnaty® XFG 10 micrograms/dose dispersion for injection COVID-19 mRNA vaccine course consists of 2 separate doses of 0.3ml each, a minimum of 3 months (12 weeks) apart.

In those who have already received one previous dose of COVID-19 vaccination a single dose of Comirnaty® XFG 10 micrograms/dose dispersion for injection COVID-19 mRNA Vaccine dose, should be given at least 3 months (12 weeks) after any previous dose of COVID-19 vaccine (regardless of the vaccine given for the previous dose) to complete the primary course.

Reinforcing dose

For those who have previously received primary vaccination a single dose of Comirnaty® XFG 10 micrograms/dose dispersion for injection COVID-19 mRNA vaccine should be given.

The dose should be given at least 3 months (12 weeks) after the previous dose of COVID-19 vaccine (regardless of the vaccine given for the previous dose).

The only exception to the 3 months interval would be where individuals were about to receive or increase the intensity of an immunosuppressive treatment, and therefore a better response would be made if immunised prior to that treatment commencing. In this unusual scenario, the interval may be reduced to a minimum of 3 weeks on specialist clinical advice.

Individuals anticipating immunosuppressive therapy

Individuals anticipating immunosuppressive therapy who have been assessed as eligible for COVID-19 vaccine and referred by their clinician for vaccination before starting treatment, should be offered vaccination in line with correspondence/publications from [Scottish Government](#). Administration is permitted under this PGD.

Individuals identified as meeting the definition for severe immunosuppression

Additional doses for those identified as meeting the definition for severe immunosuppression (as defined in [The Green Book, COVID-19 Chapter](#)) may be required.

From 2023, for most individuals aged 5 years and above, the primary course of COVID-19 vaccine is an offer of a single dose of vaccine, provided only during seasonal campaigns. Individuals who become or have recently become severely immunosuppressed (i.e. those commencing immunosuppressive therapy or those

who have developed an immunosuppressive condition) should be considered for additional doses (as outlined below).

Previously unvaccinated individuals who become or have recently become severely immunosuppressed should be considered for a first dose of vaccination, regardless of the time of year. Further doses should then be offered on the basis of specialist clinical judgement (see below).

Vaccinated individuals who become or have recently become severely immunosuppressed should be considered for an additional dose of COVID-19 vaccine, regardless of their past vaccination history and the time of year. The additional dose of vaccine should be offered at a minimum interval of 3 months from any previous doses, to extend protection until the next seasonal campaign.

Clinical judgement should be used to decide which individuals should be given an additional dose soon after their diagnosis rather than waiting for the next campaign and thus getting extra protection during the season, particularly over the winter, and at the same time as other high risk groups. The optimal timing should also take account of the degree of immune suppression (see [The Green Book COVID-19 Chapter](#) section on timing). Second doses should ideally be given between 8-12 weeks from the previous dose, to extend protection. This interval may be reduced to 3 weeks on specialist clinical advice to maximise short term protection, bearing in mind that response may be less durable. As above, subsequent doses may be optimally delivered during the next regular campaign.

In contrast to other eligible risk groups, those who are eligible for a vaccination due to severe immunosuppression but miss vaccination during the campaign period, may be considered for a booster at a later date based on individual clinical judgement, balancing their immediate level of risk against the advantages of waiting till the next seasonal campaign.

Additional doses are covered by this PGD.

Revaccination of individuals who have received a haematopoietic stem cell transplant or CAR-T treatment

In accordance with the [Revaccination of patients following haematopoietic stem cell transplant or CAR-T treatment guidance](#).

2.5. Duration of treatment

See frequency section.

2.6. Maximum or minimum treatment period

See frequency section.

2.7. Quantity to supply/administer

See frequency section.

2.8. ▼ black triangle medicines

Yes.

Healthcare professionals and individuals/carers should report suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Yellow Card reporting scheme on <http://www.mhra.gov.uk/yellowcard>.

2.9. Legal category

Prescription only medicine (POM).

2.10. Is the use outwith the SmPC?

The SmPCs for all strengths of Comirnaty® XFG COVID-19 mRNA vaccines recommend close observation for at least 15 minutes following vaccination. In recognition of the need to accelerate delivery of the programme in response to the emergence of the Omicron variant (in 2021), the UK Chief Medical Officers **recommended temporary suspension** of this requirement. This temporary suspension in individuals without a history of allergy was also agreed by the Commission on Human Medicines. The advice to suspend the routine 15 observation period has since been extended to all the currently available COVID-19 vaccines when used in the NHS programme.

For previously unvaccinated infants and children aged 6 months to 4 years, the SmPC for Comirnaty® XFG 10micrograms/dose recommends a 2 dose primary course administered 8 weeks apart. This is superseded by official recommendations for a minimum dosage interval of 3 months (12 weeks) as set out in **The Green Book COVID-19 Chapter**.

Reducing dose interval to a minimum of 3 weeks under specialist advice for individuals about to receive or increase the intensity of an immunosuppressive treatment is out with the SmPC for Spikevax® XFG but in line with official recommendations (see **The Green Book COVID-19 Chapter**).

Revaccination of patients following haematopoietic stem cell transplant or CAR-T treatment is considered off-label but is in accordance with the **Revaccination of patients following haematopoietic stem cell transplant or CAR-T treatment** guidance.

Where a vaccine is recommended off-label consider, as part of the consent process, informing the individual, parent or carer that the vaccine is being offered in accordance with national guidance but that this is outside the product licence.

Vaccine should be stored according to the conditions detailed in the Storage section below. However, in the event of an inadvertent or unavoidable deviation of these conditions refer to NHS Board guidance on storage and handling of vaccines. Where vaccine is assessed in accordance with these guidelines as appropriate for continued use this would constitute off-label administration under this PGD.

2.11. Storage requirements

General requirements

During storage it is recommended that the vaccine products are stored in the original packaging/cartons, away from direct sunlight to protect from light and kept upright.

NHS Board guidance on Storage and Handling of vaccines should be observed.

In the event of an inadvertent or unavoidable deviation of these conditions, vaccine that has been stored outside the conditions stated above should be quarantined and risk assessed for suitability of continued use or appropriate disposal.

The manufacturer may advise of updated storage requirements and product stability: as new data becomes available, vaccine should be stored in accordance with updated recommendations from the manufacturer.

Vaccine specific requirements

Spikevax® XFG 0.1mg/1ml (50 micrograms/0.5ml dose) COVID-19 mRNA

Vaccine dispersion for injection

Unpunctured

When stored for 12 months at -50°C to -15°C, once thawed and stored at 2°C to 8°C and protected from light, the unopened vial has a maximum of 14 days shelf life. This

increases to a maximum of 30 days when the vials are removed from the freezer within 9 months. 2°C to 8°C shelf life should be indicated on the outer packaging.

The unopened vaccine may be stored at 8°C to 25°C up to 24 hours after removal from refrigerated conditions.

Punctured

Chemical and physical in-use stability has been demonstrated for 6 hours at 2°C to 25°C after initial puncture. From a microbiological point of view, the product should be used immediately. If the vaccine is not used immediately, in-use storage times and conditions are the responsibility of the user.

Thawed vials can be handled in room light conditions. The vaccine vial has space to write the date and time that the vial should be discarded following first puncture; write this on the vial label.

Do not re-freeze.

Comirnaty® XFG 30 micrograms/dose dispersion for injection COVID-19 mRNA vaccine, pre-filled syringe

The vaccine should be stored refrigerated at +2°C to +8°C in the original package in order to protect from light.

Do not freeze.

Prior to use, pre-filled syringes can be stored for up to 12 hours at temperatures between 8 °C and 30 °C and can be handled in room light conditions.

Comirnaty® XFG 10 micrograms/dose dispersion for injection COVID-19 mRNA vaccine, single dose vial

The vaccine will be delivered in a thawed state. Once thawed, the unopened vial may be stored refrigerated at +2°C to +8°C protected from light for up to 10 weeks (within shelf life). Once thawed the vaccine should not be re-frozen.

Prior to use, the unopened vials can be stored for up to 12 hours at temperatures between 8 °C and 30 °C.

Thawed vials can be handled in room light conditions.

2.12. Additional information

Minor illnesses without fever or systemic upset are not valid reasons to postpone immunisation. If an individual is acutely unwell, immunisation should be postponed until they have fully recovered.

3. Adverse reactions

3.1. Warnings including possible adverse reactions and management of these

The most frequently reported adverse reactions are injection site pain, swelling or redness, fatigue, headache, fever myalgia and chills. These reactions are usually mild or moderate in intensity and resolve within a few days after vaccination.

Table 1. Summary of common and very common adverse reactions for COVID-19 vaccines

Vaccine	Very Common	Common
Comirnaty® XFG	injection site pain, tenderness and swelling, fatigue, headache, drowsiness, myalgia, arthralgia, chills, pyrexia, vomiting, diarrhoea, irritability, decreased appetite.	injection site redness, nausea, lymphadenopathy
Spikevax® XFG	lymphadenopathy, injection site pain, swelling and erythema, headache, fatigue, arthralgia, myalgia, chills, pyrexia, nausea, vomiting	injection site rash and urticaria, rash, diarrhoea

Lymphadenopathy: Swollen axilla or neck glands on the same side as the vaccination site can occur as a very common reaction, which can last for up to 10 days. If the vaccine recipient is due to attend for a mammogram, they should be advised to inform clinicians regarding date of vaccine administration.

Myocarditis and pericarditis: Very rare reports of myocarditis and pericarditis have been observed following vaccination with mRNA COVID-19 vaccines. The reported rate appears to be highest in those under 25 years of age and in males, and after the second dose. Onset is within a few days of vaccination and most cases are mild and

have recovered without any sequelae. The MHRA has advised the benefits of vaccination still outweigh any risk in most individuals. Healthcare professionals should be alert to the signs and symptoms of myocarditis and pericarditis. Recipients, and if appropriate parents and carers, should be instructed to seek immediate medical attention if they develop symptoms indicative of myocarditis or pericarditis such as (acute and persisting) chest pain, shortness of breath or palpitations following vaccination. Healthcare professionals should consult guidance and/or specialists to diagnose and treat this condition.

Heavy menstrual bleeding has been reported after COVID-19 vaccination. In most cases, this is self-limiting.

Uncommonly, benign and self-limiting cases of Erythema Multiforme have been reported associated after vaccination.

As with all vaccines there is a very small possibility of anaphylaxis and facilities for its management must be available. In the event of a severe adverse reaction individuals should be advised to seek medical advice.

For full details/information on possible adverse reaction, refer to manufacturer's product literature or SmPC.

3.2. Reporting procedure for adverse reactions

Healthcare professionals and individuals/carers should report all suspected adverse reactions to the Medicines and Healthcare products Regulatory Agency (MHRA) using the Yellow Card reporting scheme on <http://www.mhra.gov.uk/yellowcard>.

Any adverse reaction to a vaccine should be documented in accordance with locally agreed procedures in the individual's record and the individual's GP should be informed.

Anaphylaxis is a very rare, recognised side effect of most vaccines and suspected cases should be reported via the MHRA Yellow Card Scheme. **The Green Book Vaccine safety and adverse events following immunisation chapter (8)** gives detailed guidance on distinguishing between faints, panic attacks and the signs and

symptoms of anaphylaxis. If a case of suspected anaphylaxis meets the clinical features described in **The Green Book Vaccine safety and adverse events following immunisation chapter (8)**, this should be reported via the Yellow Card Scheme as a case of ‘anaphylaxis’ (or if appropriate ‘anaphylactoid reaction’). Cases of less severe allergic reactions (i.e. not including the clinical features of anaphylaxis) should not be reported as anaphylaxis but as ‘allergic reaction’.

Programmatic Adverse Events should be recorded in line with local procedures and where appropriate escalated in accordance with the national framework.

3.3. Advice to patient or carer including written information

Written information to be given to individual, parent or carer:

- Provide manufacturer’s consumer information leaflet/patient information leaflet (PIL) provided with the vaccine.
- Immunisation promotional material may be provided as appropriate

Individual advice/follow up treatment:

- Inform the individual, parent or carer of possible side effects and their management.
- Inform the individual, parent or carer that anyone who has any of the following symptoms after vaccination should seek medical advice urgently:
 - chest pain
 - shortness of breath
 - feelings of having a fast-beating, fluttering, or pounding heart
- As has always been recommended, any fever after vaccination should be monitored and if individuals are concerned about their health at any time, they should seek advice from their GP or NHS24.

- The individual should be advised to seek medical advice in the event of a severe adverse reaction.
- Inform the individual that they can report suspected adverse reactions to the MHRA using the Yellow Card reporting scheme on:
<http://www.mhra.gov.uk/yellowcard>
- As with all vaccines, immunisation may not result in protection in all individuals. The individual or carer should be advised that immunosuppressed individuals may not make a full immune response to the vaccine.
- When administration is postponed advise the individual how future vaccination may be accessed.

3.4. Observation following vaccination

Following immunisation, patients remain under observation in line with NHS Board policy.

Following COVID-19 vaccine administration, individuals should be observed for any immediate reactions whilst they are receiving any verbal post vaccination information and exiting the centre.

Syncope (fainting) can occur following vaccination. Some of the adverse effects and syncope may temporarily affect the ability to drive or use machines. It is advised not to drive or use machinery until the prevailing symptoms have cleared. Where relevant, all vaccinees should either be driven by someone else or should not drive for 15 minutes after vaccination.

An observation period when indicated after clinical assessment in individuals with a history of allergy as set out in Table 5 and flowchart in **The Green Book COVID-19 Chapter**.

Vaccinated individuals should be informed about how to access immediate healthcare advice in the event of displaying any symptoms. In some settings, for

example domiciliary vaccination, this may require a responsible adult to be present for at least 15 minutes after vaccination.

3.5. Follow up

As above.

3.6. Additional facilities

A protocol for the management of anaphylaxis and an anaphylaxis pack must always be available whenever vaccines are given. Immediate treatment should include early treatment with intramuscular adrenaline, with an early call for help and further IM adrenaline every 5 minutes.

The health professionals overseeing the immunisation service must be trained to recognise an anaphylactic reaction and be familiar with techniques for resuscitation of a patient with anaphylaxis.

4. Characteristics of staff authorised under the PGD

4.1. Professional qualifications

The following classes of registered healthcare practitioners are permitted to administer vaccines included in this PGD:

- nurses and midwives currently registered with the Nursing and Midwifery Council (NMC).
- pharmacists currently registered with the General Pharmaceutical Council (GPhC).
- pharmacy technicians currently registered with the General Pharmaceutical Council (GPhC).
- chiropodists/podiatrists, dieticians, occupational therapists, orthoptists, orthotists/prosthetists, paramedics, physiotherapists, radiographers and speech and language therapists currently registered with the Health and Care Professions Council (HCPC).
- dental hygienists and dental therapists registered with the General Dental Council.
- optometrists registered with the General Optical Council.

4.2. Specialist competencies or qualifications

Persons must only work under this PGD where they are competent to do so.

All practitioners operating this PGD:

- demonstrate appropriate knowledge and skills to work under the PGD for the administration of COVID-19 vaccine.

- have met the requirements of the Public Services Delivery Scotland (PSDS) **Proficiency Document for Administration of Vaccines by Registered Healthcare Practitioners** and **COVID-19 vaccination core eLearning**.

All persons operating this PGD:

- must be authorised by name by their employer as an approved person under the current terms of this PGD before working to it
- must be familiar with the vaccine products and alert to changes in the manufacturer's product information/summary of product characteristics information
- must be competent to undertake immunisation and to discuss issues related to immunisation to assess patients for vaccination and obtain consent
- must be competent in the correct storage of vaccines and management of the cold chain if receiving, responsible for, or handling the vaccine
- must be competent in the recognition and management of anaphylaxis or under the supervision of persons able to respond appropriately to immediate adverse reactions
- must have access to the PGD and associated online resources
- should fulfil any additional requirements defined by local policy.

All practitioners operating under the PGD are responsible for ensuring they remain up to date with the use of COVID-19 vaccines included. If any training needs are identified these should be discussed with the individuals in the organisation responsible for authorising individuals to act under the PGD.

Employer

- The employer is responsible for ensuring that persons have the required knowledge and skills to safely deliver the activity they are employed to provide under this PGD.
- As a minimum, competence requirements stipulated in the PGD must be adhered to.

4.3. Continuing education and training

All practitioners operating under the PGD are responsible for ensuring they remain up to date with the use of vaccines included. If any training needs are identified these should be discussed with the individuals in the organisation responsible for authorising individuals to act under this PGD.

5. Audit trail

Record the following information:

- valid informed consent was given
- name of individual, address, date of birth and GP with whom the individual is registered if possible
- name of person that undertook assessment of individual's clinical suitability and subsequently administered the vaccine
- name and brand of vaccine
- date of administration
- dose, form and route of administration of vaccine
- batch number
- where possible expiry date
- anatomical site of vaccination
- advice given, including advice given if excluded or declines immunisation
- details of any adverse drug reactions and actions taken
- administered under PGD

Records should be kept in line with local procedures.

Local policy should be followed to encourage information sharing with the individual's General Practice.

All records should be clear, legible and contemporaneous and in an easily retrievable format.

6. Additional references

Practitioners operating the PGD must be familiar with:

- **Immunisation against Infectious Disease [Green Book]**
- **Immunisation against Infectious Disease [Green Book] COVID-19**
- **Spikevax® XFG 0.1 mg/mL COVID-19 mRNA vaccine Summary of Product Characteristics**
- **Comirnaty® XFG 30 micrograms/dose COVID-19 mRNA vaccine Summary of Product Characteristics**
- **Comirnaty® XFG 10 micrograms/dose COVID-19 mRNA vaccine Summary of Product Characteristics**
- **Educational resources for registered professionals produced by Public Services Delivery Scotland**
- **All relevant JCVI statements**
- **All relevant Scottish Government advice including the relevant CMO letter(s)**
- **Revaccination of patients following haematopoietic stem cell transplant or CAR-T treatment schedule**
- **Prescribing and dispensing / supply / administration by the same healthcare professional - Royal College of Pharmacy.**
- **Professional guidance to support prescribing and dispensing / supply / administration by the same healthcare professional - Royal College of Pharmacy.**
- **Scottish Government Section 47 certificate of incapacity**
- **Public Services Delivery Scotland: Adults with incapacity**

- **COVID-19 vaccination programme: information for healthcare practitioners - GOV.UK**

7. PGD for administration of COVID-19 mRNA vaccine to individuals from 6 months of age version 1.0 (valid from 24 September 2026 and expires 30 April 2027): authorisation

Practitioner

This PGD does not remove professional obligations and accountability. It is the responsibility of each professional to practice within the bounds of their own competence and in accordance with their Code of Professional Conduct and to ensure familiarity with the marketing authorisation holder's summary of product characteristics for all vaccines administered in accordance with this PGD.

By signing this Patient Group Direction, you are indicating that you agree to its contents and that you will work within it. I agree to administer COVID-19 vaccine to individuals from 6 months of age and over only in accordance with this PGD.

Name of professional	Signature	Date

Authorising manager

I confirm that the practitioners named above have declared themselves suitably trained and competent to work under this PGD. I give authorisation on behalf of **NHS Greater Glasgow and Clyde** for the above-named healthcare professionals who have signed the PGD to work under it.

Lead clinician for the service area:

Name

Signature

Date

Authorised staff should be provided with an individual copy of the clinical content of the PGD and a photocopy of the document showing their authorisation.

This authorisation sheet should be retained to serve as a record of those practitioners authorised to work under this PGD.

8. Version history

Version	Date	Summary of changes
1.0	24 September 2026	New combined PGD template to support delivery of the Autumn 2026 COVID-19 vaccination programme to eligible individuals aged 6 months and over. This PGD reflects the change in antigenic content of the COVID-19 vaccine, from LP.8.1 (as utilised in Spring 2026 campaign) to XFG. The PGD also amalgamates the 2 previously separate PGDs for adults and children aged under 18 years, into a single legal framework.