

Bacterial Respiratory Infections Service

SMiRL (Glasgow)
Level 5, New Lister Building
Greater Glasgow Glasgow Royal Infirmary,
and Clyde
10-16 Alexandra Parade,
Glasgow G31 2ER
0141 242 9633

BRIS Request Form

Do you suspect that any of the isolates/specimens you are referring could be Hazard Group 3 or 4? ☐ Yes ☐ No Please provide further details/preliminary ID results below.

** SMiRL USE ONLY **					
SMiRL code					
Booked in by					
Checked by					
Scan 1					
PID					
Cultured by					

PATIENT/SPECIMEN DETAILS							
CHI Number:			Sex:		Male		Female
Surname:			Address:				
Forename:							
Date of Birth:			Post Code:				
SENDER'S INFORMATION/CONTACT DETAILS							
Sending Lab/Consultant:		Sending Lab Address:					
Secondary Location (Hospital/Ward):							
Contact Number:							
SPECIMEN DETAILS							
Date/Time Collected:			Sender's Reference Number:				
Specimen Type:							
Clinical details:							
SENDING LAB RESULTS - please provide an organism ID and any relevant antibiotic MICs as per referral criteria							
Organism ID: MIC:		MIC:					
Referral Criteria - please select							
Organism	Referral Criteria						
Haemophilus influenzae	 □ Sterile site isolate (ID, AST and typing) □ All 3rd/4th & 5th gen. cephalosporin and carbapenem resistant isolates (ID, AST and typing) □ Patient sample for bacterial PCR 						
Legionella species	 □ Urine for urinary antigen ELISA □ Patient and related environmental isolates (ID and typing) □ Patient respiratory sample for PCR and culture 						
Streptococcus pneumoniae	 □ Sterile site isolate (ID, AST and typing) □ All isolates penicillin MIC > 2mg/L □ All third generation cephalosporin intermediate or resistant isolates □ All vancomycin, teicoplanin, linezolid, or rifampicin resistant isolates □ Patient sample for bacterial PCR 						
Neisseria meningitidis	 Sterile site isolate (ID, AST and typing) Non sterile site isolate from suspected cases (ID, AST and typing) Patient sample for bacterial PCR 						
Streptococcus pyogenes (GAS) Bordetella pertussis	 □ Sterile site isolate □ Necrotising fasciitis □ Serious soft tissue infection (emm typing) □ Any isolates linked to outbreaks including health care and maternity settings □ All penicillin, cephalosporin, vancomycin, teicoplanin, daptomycin, linezolid or tigecycline resistant isolates □ Serum for IgG determination to pertussis toxin 						
The second of th							

RF-4 v3

Page 1 of 1