

**CDI Assessment –
TO BE COMPLETED BY MEDICAL STAFF**



Completed by:	Date:	
CDI Severity Markers	Yes	No
Temperature >38.5°C		
Suspicion of PMC, toxic megacolon, ileus		
Colonic dilatation in CT scan/ abdominal x-ray >6cm		
WBC > 15cells/mm ³		
Creatinine >1.5 x baseline		
Total		

If two or more of the above markers are positive, this **may** indicate severe disease. Please follow AMT Treatment Algorithm and review if patient is still passing loose stools after 5 days of initial treatment.

**CDI Assessment –
TO BE COMPLETED BY MEDICAL STAFF**



Completed by:	Date:	
CDI Severity Markers	Yes	No
Temperature >38.5°C		
Suspicion of PMC, toxic megacolon, ileus		
Colonic dilatation in CT scan/ abdominal x-ray >6cm		
WBC > 15 cells/mm ³		
Creatinine >1.5 x baseline		
Total		

If two or more of the above markers are positive, this **may** indicate severe disease. Please follow AMT Treatment Algorithm and review if patient is still passing loose stools after 5 days of initial treatment.

**CDI Assessment –
TO BE COMPLETED BY MEDICAL STAFF**



Completed by:	Date:	
CDI Severity Markers	Yes	No
Temperature >38.5°C		
Suspicion of PMC, toxic megacolon, ileus		
Colonic dilatation in CT scan/ abdominal x-ray >6cm		
WBC > 15 cells/mm ³		
Creatinine >1.5 x baseline		
Total		

If two or more of the above markers are positive, this **may** indicate severe disease. Please follow AMT Treatment Algorithm and review if patient is still passing loose stools after 5 days of initial treatment.

**CDI Assessment –
TO BE COMPLETED BY MEDICAL STAFF**



Completed by:	Date:	
CDI Severity Markers	Yes	No
Temperature >38.5°C		
Suspicion of PMC, toxic megacolon, ileus		
Colonic dilatation in CT scan/ abdominal x-ray >6cm		
WBC > 15 cells/mm ³		
Creatinine >1.5 x baseline		
Total		

If two or more of the above markers are positive, this **may** indicate severe disease. Please follow AMT Treatment Algorithm and review if patient is still passing loose stools after 5 days of initial treatment.