Infection Prevention and Control Care Checklist – CPE

This Care checklist should be used with patients who are suspected of having CPE and are undergoing screening. It should also be used for those patients who have ever been /are positive during their stay in hospital and then signed off at discharge. Each criteria should be ticked V if in place or X if not, the checklist should be then initialled after completion, daily.

Date Isolation commenced:

Patient Name:

CHI:

	Patient Placement/ Assessment of Risk	Daily check (v/x)					
Patient Placement /Assessment of risk	Patient isolated in a single room with en suite facilities / own commode. If						
	a single room is not available, an IPCT risk assessment is completed (see						
	Appendix 1) Stop isolation if patient has 3 negative rectal screens (and						
	never been positive) or on discharge.						
	Place yellow isolation sign on the door to the isolation room						
	Door to isolation room is closed when not in use. If for any reason this is						
	not appropriate then an IPCT risk assessment is completed.						
	Hand Hygiene (HH)				<u> </u>		
	All staff must use correct 6 step technique for hand hygiene at 5 key						
	moments						
	HH facilities are offered to patient after using the toilet and prior to						
	mealtimes etc. (Liquid soap and water/wipes where applicable)						
	Personal Protective Clothing (PPE)				<u> </u>		
us	Disposable gloves and yellow apron are worn for all direct contact with						
tio	the patient and their equipment/environment, removed before leaving						
cau	the isolation area and discarded as clinical waste. HH must follow						
re	removal of PPE.						
b b	Safe Management of Care Equipment						
ase	Single-use items are used where possible OR equipment is dedicated to						
n B	patient while in isolation.						
sio	There are no non-essential items in room. (eg. Excessive patient						
mis	belongings)						
SUE	Twice daily decontamination of the patient equipment by HCW is in place						
Tra	using 1,000 ppm solution of chlorine based detergent						
8	Safe Management of Care Environment						
Standard Infection Control & Transmission Based Precautions	Twice daily clean of isolation room is completed by Domestic services,						
	using a solution of 1,000 ppm chlorine based detergent. A terminal clean						
	will be arranged on day of discharge/end of isolation.						
	Laundry and Clinical/Healthcare waste						
	All laundry is placed in a water soluble bag, then into a clear plastic bag						
	(brown bag used in mental health areas), tied then into a red laundry bag.						
	Clean linen must not be stored in the isolation room.						
	All waste should be disposed of in the isolation room as clinical/						
	healthcare waste.						
	Information for patients and their carers						
ers	The patient has been given information on CPE. This is available as a						
nts/carers	narrated video via a QR code or a paper copy of the Patient Information						
ts/e	Leaflet (PIL) is available in several languages.						
oati							
or							
5	<u> </u>						
Information for patie	If taking clothing home, carers have been issued with a Washing Clothes						
Ĕ	at Home patient information leaflet (PIL).						
lfo	(NB. Personal laundry is placed into a domestic water soluble bag, then into a patient clothing bag before being given to carer to take home)						
-							
	HCW Daily Initial :			Γ			

Date Isolation ceased/ Terminal Clean completed:

Care Checklist completed and signed off by:

Appendix 1: Infection Prevention and Control Risk Assessment			Addressograph Label:					HS	
(for patients with known or suspected infection that cannot be	e isolated)	olated) Patient Name and DOB/CHI:				1:	Greater Glasgow and Clyde		
Daily Assessment / Review Required									
	C O M M E N T S	DATE	DATE	DATE	DATE	DATE	DATE	DATE	
Daily Assessment Performed by Initials									
Known or suspected Infection e.g. unexplained loose stools, MRSA, Group A Strep, <i>C. difficile</i> , Influenza, pulmonary tuberculosis. <i>Please state</i>									
Infection Control Risk , e.g. unable to isolate, unable to close door of isolation room.									
Please state									
Reason unable to isolate / close door to isolation room, e.g. falls risk, observation required, clinical condition.									
Please state									
Additional Precautions put in place to reduce risk of transmission, e.g. nursed next to a clinical wash hand basin, at end of ward, trolley containing appropriate PPE at end of bed, next to low risk patient, clinical waste bin placed next to bed space. <i>Please state</i>									
Infection Prevention and Control have been informed of patient's admission and are aware of inability to adhere to IPC Policy?									
Yes / No									
Summary Detail of Resolution									
Daily risk assessments are no longer required		Signed Date	k						