

## <u>CPE – Infection Prevention and Control</u> Infection Prevention and Control Advice

Patient: has	S CPE (Carbapenemase producing
enterobacteraeicae)	
Isolated in a specimen taken on the//	'
The patient should be isolated in a single r	oom with standard and transmission based
precautions, for the duration of their stay	in hospital. The following should be

implemented:

- the patient room /bed space and equipment should be cleaned twice per

- day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed.
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

IPCN:	Date	1	/
IPCN:	Date:	//	



## <u>CPE – Infection Prevention and Control</u> Infection Prevention and Control Advice

Patient:	has CPE (Carbapenemase producing
enterobacteraeicae)	

Isolated in a specimen taken on the --/---/---.

The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- the patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed.
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

IPCN:	Date:	/	/
II CIVI	Date.	•••/ •••	,



## <u>CPE – Infection Prevention and Control</u> Infection Prevention and Control Advice

Patient:	has CPE (Carbapenemase producing
enterobacteraeicae)	

Isolated in a specimen taken on the --/---/---.

The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- the patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed.
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

IPCN:	Date:	//	/
II VIV	Dutc.	//	• •



## <u>CPE – Infection Prevention and Control</u> <u>Infection Prevention and Control Advice</u>

Patient:	has CPE (Carbapenemase producing
enterobacteraeicae)	

Isolated in a specimen taken on the --/---/--- .

The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- the patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed.
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

PCN: Date:	//		
Ducc.	• • • • • •	• • • • •	•