

**Patient:** .....has CPE (*Carbapenemase producing enterobacteraeiae*)  
Isolated in a specimen taken on the --/--/--- .  
The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed .
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

**IPCN:** ..... **Date:** .../.../.....

**Patient:** .....has CPE (*Carbapenemase producing enterobacteraeiae*)  
Isolated in a specimen taken on the --/--/--- .  
The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed .
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

**IPCN:** ..... **Date:** .../.../.....

**Patient:** .....has CPE (*Carbapenemase producing enterobacteraeiae*)  
Isolated in a specimen taken on the --/--/--- .  
The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed .
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

**IPCN:** ..... **Date:** .../.../.....

**Patient:** .....has CPE (*Carbapenemase producing enterobacteraeiae*)  
Isolated in a specimen taken on the --/--/--- .  
The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed .
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

**IPCN:** ..... **Date:** .../.../.....

**Patient:** .....has CPE (*Carbapenemase producing enterobacteraeiae*)  
Isolated in a specimen taken on the --/--/--- .  
The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- the patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed .
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

**IPCN:** ..... **Date:** .../.../.....

**Patient:** .....has CPE (*Carbapenemase producing enterobacteraeiae*)  
Isolated in a specimen taken on the --/--/--- .  
The patient should be isolated in a single room with standard and transmission based precautions, for the duration of their stay in hospital. The following should be implemented:

- The patient room /bed space and equipment should be cleaned twice per day with chlorine based detergent
- a disposable apron and gloves should be used when delivering care directly.
- the door to the room should be closed .
- the CPE IPC care checklist should be completed daily.
- Further screening should be discussed with the local IPCT
- The IPCT will place a CPE tag on patients notes in TrakCare

**IPCN:** ..... **Date:** .../.../.....

