



# TITAN CHLOR PLUS® GENERAL ENVIRONMENT

WHEN URINE, VOMIT, FAECES ARE PRESENT OR IF INFECTION IS KNOWN OR SUSPECTED



Wear disposable gloves and aprons. Safety glasses must be worn whilst handling the chemical solution



Remove any gross contamination including urine, vomit (do not use directly on urine and vomit) and faeces with disposable paper before applying TITAN Chlor Plus. Dispose of as healthcare waste.



To make solution, dissolve TITAN Chlor Plus tablets (according to dilution instructions below) in cold water. Leave lid off dilution bottle until the tablets have dissolved. Mark date and time prepared on dilution bottle.



Use solution according to hospital policies. When disinfecting items such as commodes and mattresses, allow to dry fully before rinsing.



Keep solution for no longer than 24 hours. Dispose of remaining solution into DSR or sluice sink drain (not healthcare hand wash sinks). Rinse with running water.



Dispose of gloves and apron as clinical waste. Wash hands after removing gloves and apron.

## DILUTION INSTRUCTIONS

### DISINFECTION OF THE ENVIRONMENT (1,000 PPM AVAILABLE CHLORINE)

### CONTACT TIME

TITAN Chlor Plus Tablets

1 Tablet per 1 L cold water

Disinfection of the environment

5 minutes

Sporicidal - C.difficile

5 minutes

## WARNINGS & PRECAUTIONS



**AVOID** contact with skin and eyes.  
**DO NOT** breathe dust/fume/gas/mist/vapors/spray.



**DO NOT** use directly on urine or vomit. Mixing this product with acid or ammonia releases chlorine gas.



**AVOID** prolonged contact with stainless steel or clothing.



**ALWAYS** dispose used materials as clinical waste. Wash hands thoroughly after handling.



**ALWAYS** keep container tightly closed. Store in a secure dry place in suitable labelled containers.



**ALWAYS** keep out of reach of children.



**ALWAYS** dispose of each solution after 24 hours. Refer to your SDS and hospital policy.



**USE ONLY** with adequate ventilation.