


NHSGGC Alcohol and Drug Recovery Services **Cocaine Toolkit**



Lead/Authors:	MAT SPMG
Document Type:	Guidance
Approved By:	Medicated Assisted Treatment – Substitute Prescribing Management Group
Date:	V1- 24.05.2022 V2 Review- 01.09.2025
Version:	V2
Review Date:	2 years from review

Contents

Working with people at risk of harm from cocaine use - Guidance for ADRS front line staff...	5
Executive Summary.....	5
Foreword	5
Acknowledgements	6
Introduction.....	7
Whole Family Approach	8
Assessing cocaine use and impact.....	9
Guidance notes.....	9
An overview of Cocaine	9
Forms of cocaine and associated cost	12
The local powder cocaine market	12
Cocaine and other drugs.....	14
Cocaine and alcohol.....	14
Cocaine and heroin	14
Cocaine and benzodiazepines	14
Cocaine and ketamine	15
Cocaine compulsion and psychological dependence.....	16
Stages of cocaine withdrawal and recovery	17
Cocaine harms and harm reduction	18
Cocaine Overdose Stroke Heart attack Toxicity (COSHT).....	18
Signs and Symptoms of COSHT.....	19
Clinical Outcomes	19
Reducing the Risk of COSHT.....	20
Cocaine and sexual risks.....	21
Cocaine, Malnutrition and the Immune System.....	22
Supporting Better Nutrition and Self Care.....	22
Risks associated with methods of administration	24
Snort risks	24
Transmission of infections including blood borne viruses.....	25
Snort harm reduction.....	25
Smoke risks	25
Pipe risks.....	26
Recycling cocaine from pipes	27
Smoke harm reduction	27
Injecting related harm	28
Blood Borne Virus (BBV) risks.....	28
Injecting related complications and injury.....	28
Preparing cocaine for injection.....	30
Cocaine injecting harm reduction	31
Needle and Syringe Guide.....	32

Physical health monitoring and treatment interventions for individuals who are prescribed OST and are using cocaine.....	33
Mental health effects and treatment interventions	34
Crisis intervention	34
Symptoms of drug related mental health crisis.....	34
Common symptoms of drug related mental health crisis.....	34
Responding to a drug induced mental health crisis	35
De-escalation	35
Five things grounding technique.....	36
Next steps.....	37
Mental Health Assessment	37
Psycho-social Interventions.....	39
Stage 1: Motivational Enhancement	39
Stage 2: Developing control over impulsive behaviour	39
Identifying and managing triggers	40
Coping with cravings	40
Seemingly irrelevant decisions and high risk situations	40
Stage 3: Developing a rewarding lifestyle	41
Managing complex cases	42
Bibliography	43
Appendices	44
Cocaine Assessment Tool	44
Motivational Enhancement: Identifying Problems	56
Motivational Enhancement: Identifying Things to Work On	59
Motivational Enhancement: Daily Routine Worksheet	62
Motivational Enhancement: Happiness Scale	64
Motivational Enhancement: Exploring the Relationships between Substance Use and Mental Health	65
Motivational Enhancement: What I Want From Treatment:	66
Motivational Enhancement: Exploring the Pros and Cons of Making a Change	70
Developing Control: Functional Analysis	71
Developing Control: Functional Analysis of Recovery-Orientated Behaviour	72
Developing Control: Discovering Triggers	73
Discovering Triggers of Your Alcohol and Other drugs Use.....	73
Developing Control: Self-Management Plan	76
Developing Control: Coping with Cravings and Urges	77
Developing Control: Coping with Thoughts about Using	79
Developing Control: Problem-solving Worksheet	80
Developing Control: Seemingly Irrelevant Decisions	82
Developing a Rewarding Lifestyle: Recreational Survey	83
Developing a Rewarding Lifestyle: Brief Goal Setting Worksheet for Reinforcing Recovery	87
Developing a Rewarding Lifestyle: Highlighting Progress Graph	88
Developing a Rewarding Lifestyle: Highlighting Progress Calendar	89
Developing a Rewarding Lifestyle: Timeline Exercise.....	90

Working with people at risk of harm from cocaine use - Guidance for ADRS front line staff

Executive Summary

Data from over 5000 comprehensive Assessments of Injecting Risk (AIR) suggests that cocaine has become the primary drug of injection for most people who prepare and inject drugs in Greater Glasgow and Clyde.

Although we now have cocaine only injecting groups, there are people who also inject heroin or alternate between the two. Many individuals who inject cocaine and heroin are linked with services and receiving Opioid Substitution Therapy (OST). This provides us with an opportunity to explore their cocaine use and suggest a plan to reduce harm and facilitate treatment. We must do this in order to support individuals holistically and address the real harms that they face.

Many of the harm reduction techniques and treatment options available to people using cocaine vary greatly from the traditional interventions we use when working with people using opioids. There is currently no substitute prescribing option for people using cocaine. For those at risk of overdose from cocaine, there is no equivalent drug to naloxone, which is used in opioid overdose to quickly reverse symptoms. Advice on preparation and methods of administration of cocaine vary, as do the implications of polydrug and alcohol use when combined with cocaine. Therefore, it is vitally important that through training we develop skills and become familiar with the approaches needed to engage with individuals using cocaine.

Similarly, the presentation of people using cocaine and stimulants can be quite different to those using depressant drugs alone. An understanding of these differences in presentation can be key to identifying the best supports for people. In particular, cocaine has a significant impact on mental and physical health, both acutely and over a period of time. The sections on mental health and physical health within the toolkit will give the reader a greater understanding of these risks and the skills to use to identify and address them.

This toolkit provides educational content on cocaine and its impact on the person taking the drug, along with current detail of street terms used, costs and the local drug market. It also offers practical tools to use directly with individuals, such as the assessment tool, advice on crisis intervention and de-escalation, along with the motivational enhancement, developing control and developing a rewarding lifestyle worksheets.

We would encourage anyone working with people who use cocaine to navigate through the document in their own time and print or share resources to support those using cocaine, those impacted by cocaine use and also other allied services or service providers.

Foreword

The development of the Working with People at Risk of Harm from Cocaine Use Guidance is part of the work plan of the Medicated Assisted Treatment – Substitute Prescribing Management Group (MAT SPMG), NHS GG&C Alcohol and Drug Recovery Services (ADRS) Care Governance Subgroup.

Members involved in the development of the guidance comprised of experts from a range of disciplines. These included doctors, pharmacists and psychologists from ADRS and representatives from residential services, Injecting Equipment Provision (IEP) Services and the Scottish Drugs Forum (SDF) Peer Research and Engagement Services.

A key stage in the development of the guidance involved peer volunteers and people with lived or living experience to assist in identifying the particular areas to focus on. By involving individuals with peer, lived and living experience it allowed for a greater understanding of cocaine use and associated issues. This voluntary involvement was greatly appreciated and a crucial stage in the development of the guidance.

Acknowledgements

In acknowledging the many people who have contributed directly and indirectly to this work, we would particularly like to thank the following:

Dr Katie Browne (Senior Medical Officer, ADRS)
Dr Beate Beck-Schwahn (Consultant Psychiatrist, ADRS)
Dr Peter Brennan (Principal Clinical Psychologist, ADRS)
John Campbell (IEP Improvement Manager, ADRS)
Dr Anna Fletcher (Consultant Addiction Psychiatrist, ADRS)
Stephanie Hartley (Cognitive Behavioural Therapist, ADRS)
Amanda King (Research & Care Governance Assistant, ADRS)
Katy MacLeod (Peer Research & Inclusion Programme Manager, SDF)
Karen Rowell (Advanced Pharmacist Independent Prescriber, GADRS)

Copyright © NHS Greater Glasgow and Clyde 2025  CC BY-NC-SA 4.0 unless otherwise noted.

Introduction

Individuals and services are reporting more poly drug use involving cocaine than ever before. Cocaine was implicated in 41% of drug related deaths in Scotland in 2023. Data from the ADRS WAND (Wound care, Assessment of Injecting, Naloxone & Dried Blood Spot Testing) Initiative in NHS GG&C between April-March 2024/25 shows that the number of people injecting cocaine is higher than those injecting heroin (84% cocaine versus 51% heroin). Of those who report injecting heroin or cocaine, 50% also report the smoking of crack or freebase cocaine. With this comes an additional range of associated harms.

Unlike opioid substitution treatment (OST) for people using heroin, there is no replacement medication available for people using cocaine, although research is continuing to explore possible treatments. Psycho-social interventions remain the cornerstone of treatment.

Below are some basic principles which may be helpful to guide front line staff working with people at harm from cocaine use. The guidance is primarily written for staff working with people currently in treatment with OST for opiate dependence and using cocaine, but may also be relevant to individuals presenting with primary cocaine use.

Whole Family Approach

Drug and Alcohol Services are underpinned by family inclusive practice. People will be given the opportunity to involve family in their care or express their wishes not to have information shared. Services have a role in supporting families and signposting them to partner organisations to access additional support where required. Staff will consider if people have caring responsibilities and offer support with this.

COCAINE AND KEEPING CHILDREN SAFE

Cocaine use in pregnancy

Increases risks of:

- Miscarriage
- Early labour
- Babies dying suddenly in their sleep



Cocaine harm to children

- Babies can have cocaine withdrawal symptoms
- Cocaine can get into breast milk
- Babies and children can be exposed to cocaine by breathing in the drug or by touching things that have cocaine on them

Cocaine presents a risk to infants and children when used in the environment where a child is. This could be due to passive inhalation or touching surfaces contaminated with residue from cocaine use. Staff should discuss preventative measures such as handwashing, awareness of food preparation and minimising consumption in places where adults interact with children. It is essential that staff promote safe sleep guidance with patients using cocaine and other substances, including alcohol, and avoid co-sleeping. The magnitude of this risk is not yet known but may be significant.

Drug and Alcohol Services have an important role in the protection of unborn babies, children and young people. If staff believe that a child or young person is a risk of harm then NHSGGC Child Protection processes must be followed and a notification of concern submitted. Robust communication with other agencies involved in the care of the family is key.

The Public Protection Service can provide advice and support to any staff who have a concern about a child or young person. The Public Protection Service in NHS GG&C can be contacted on **0141 451 6605**.

Assessing cocaine use and impact

Guidance notes

This is an assessment tool that can be used with people using cocaine, with a particular focus on injecting cocaine and smoking crack/freebase cocaine. It is not intended for use with people who use cocaine on a more occasional or recreational basis. The tool should be used to better understand an individual's cocaine use, including the patterns of use and potential risks and harms. It is intended to help highlight areas where further exploration and support may be required. It does not replace a formal physical, mental health or sexual health assessment.


The assessment does not have to be completed in any particular order and works best when completed in conversation with the person using cocaine. The questions act as prompts to open up discussion, rather than requiring staff to read all questions verbatim. This allows focus on the relevant questions for the person and ensures the full assessment can be completed comfortably within one appointment.

It is recognised that for some people it may not be possible to complete the assessment in its entirety, or that it may require to be done over more than one appointment. The more information gathered the better, and where it has been necessary to complete the assessment over several appointments, it is important to note that responses can change over time. It may be necessary to review previous answers. [Cocaine Assessment Tool](#)

An overview of Cocaine

Cocaine hydrochloride is the most common form of cocaine used in the UK. It is isolated from the leaves of the Andean coca shrub using a range of chemicals and refining processes. This plant can only be found in very specific regions of South America, meaning all the cocaine found in the UK has been trafficked here. Of course, this also means there are multiple opportunities for it to be cut, adulterated or "bashed" throughout its journey. Common agents used to cut cocaine include glucose, creatine, caffeine and benzocaine. Analysis has also shown that on occasion cocaine has been cut with painkillers such as phenacetin. Cocaine powder may be snorted or injected. Its form can also be changed to facilitate smoking.

COCAINE PRODUCTION



Regardless of how cocaine is administered, it significantly impacts neurotransmitter systems in the limbic (reward) part of the brain. It primarily affects dopamine, norepinephrine and serotonin.

Cocaine inhibits the reuptake of these neurotransmitters by binding to their transporters:

Dopamine Transporter - Binding to this transporter blocks dopamine reuptake and so increases its concentration in the synapse.

Result - Euphoria, increased energy and heightened alertness.

Norepinephrine Transporter - Binding to this transporter blocks norepinephrine reuptake and so prolongs its activity.

Result - Increased heart rate, blood pressure and alertness.

Serotonin Transporter - Binding to this transporter blocks serotonin reuptake, although this effect is less pronounced than with dopamine and norepinephrine.


Result - May contribute to initial mood elevation but plays a lesser role when compared to dopamine.

It is primarily the excess of dopamine that produces feelings of euphoria, well-being, enhanced alertness and a desire for social bonding. This is likely the main driver for compulsion to take more.

This mechanism of dopamine transporter blockade, and increased dopamine release caused by cocaine, followed by a period of dopamine depletion, creates two clear stages. Initially, a high caused by increased dopamine, followed by a state of dopamine depletion after use. Many of the negative psychological effects people experience relate to this dopamine depleted state after using.

Cocaine blocking the dopamine transporter

Cocaine



Common cocaine street terms

- Gear, coke, charlie, chico, ching and snow refer to powder cocaine
- Gram, "eggs and ham" and a 'G' all refer to the most commonly sold weight of powder cocaine which is 1 gram
- Line, patsy and patsy cline refer to a line of cocaine which is commonly snorted
- Bump or key refer to a small mound of powder cocaine which will be snorted without a straw
- Rock, base, freebase and nugget refer to crack or freebase cocaine
- Pipe and crack pipe refer to the utensil needed for smoking
- Snowball refers to mixing heroin and cocaine together for injecting

Forms of cocaine and associated cost

There are 3 main forms of cocaine used in the UK: powder, crack and freebase. Each form lends themselves to different methods of administration.

Powder Cocaine (cocaine hydrochloride) is the most common form of cocaine to be sold. It is often snorted in lines through a straw or banknote, however it is also water soluble so easy to prepare for injection. This type of cocaine does not lend itself well to smoking due to the high melting point.



The local powder cocaine market

Over the years the powder cocaine market has developed into a tiered model with ranges of purities linked to price. In Glasgow this is particularly well established with all purity levels indicated by name.

- Small bags which are aimed at those injecting are £10-£15
- Poor quality - this is often called "council" is approximately £30 per gram
- Medium quality/purity - this is often called 50-50 and costs approximately £50 per gram
- High quality/purity - this is often called "proper" or "prop" and costs approximately £80 per gram

This business model is reflective of a well-established, flexible and resilient cocaine market.

Crack Cocaine is a base form of cocaine that can be produced by mixing powder cocaine with water and sodium bicarbonate (baking soda). This is then be heated to form a solid rock like state for smoking. Although produced for smoking, these rocks can also be injected, however an acidifier such as vitamin C needs to be added to return it to a water-soluble state first. It is not uncommon for suppliers to add bulking agents during preparation to increase profit which will result in a poor-quality product – **not all crack cocaine is high purity.**



Cost - £10 - £20 per rock depending on weight and purity.

Freebase Cocaine is a base form of cocaine. The process of freeing the cocaine base comes from the adding of ammonia. The result is a form of cocaine that is very pure. This form has a low melting point which makes it easy to smoke. Freebase cocaine is not soluble in water and so an acidifier, such as vitamin C, needs to be added to return it to a water-soluble state - **all freebase cocaine is high purity.**

Cost - Not commonly sold at street level but when it is cost will always be related to weight e.g. 0.2 grams

Cocaine and other drugs

Cocaine and alcohol

The interaction between cocaine and alcohol is complex. One of the reported benefits of taking cocaine and alcohol together is the ability to consume far more alcohol without passing out or appearing very drunk. Of course, the damage caused by consuming large amounts of alcohol in a binge fashion does not disappear. This level of alcohol consumption will make people's "come down" far worse than using cocaine on its own. When both drugs are consumed together the body produces a third chemical called cocaethylene. This is known to enhance the euphoric effect of cocaine and increase the duration of effect. This may explain why so many people have such a strong urge to use both together. As well as the risks associated with increased alcohol consumption, cocaethylene creates other risks because it is toxic to the cardiovascular system. Cocaethylene elevates heart rate and blood pressure more than cocaine would alone.

Cocaine and heroin

The most common way to take heroin and cocaine together is through an injection called a "snowball". There are however several other ways the drugs can be taken together such as; injecting heroin and smoking crack/freebase, or smoking heroin and injecting cocaine. It is a commonly held false belief that when taking both heroin and cocaine at the same time they will cancel out the effects of each other. In actual fact, a synergy occurs where the effects of both drugs are significantly amplified. Overdose risk can be increased due to a false sense of relative sobriety, which is driven by cocaine and may lead to frequent re-dosing of heroin, thus increasing the risk of overdose.

Cocaine and benzodiazepines

Cocaine can be taken with benzodiazepines in an attempt to counteract some of the more sedative effects of benzodiazepines. People may also take cocaine and benzodiazepines in order to enhance, or prolong, a cocaine driven euphoric high. It is also common for people to take benzodiazepines after a cocaine binge to help ease the comedown or help sleep. There is a risk of overdose with either drug used alone, which may be increased when taken in combination. Taking these drugs together may also cause the person to feel that they can consume more which increases the likelihood of overdose.

Cocaine and ketamine

When cocaine and ketamine are taken together they can create a dreamy euphoric state. People can also feel that they have increased energy. These drugs used in combination can cause a series of life-threatening conditions. Some of the adverse effects of this combination include; unpredictable behaviour, bad trips, psychotic reactions, heart problems, high blood pressure, stroke and other neurological complications. As this market has developed, there is now evidence that these drugs are being sold 'premixed' for ease of use. In this scenario the buyer will not be aware of purity or concentrations of either drug. The street name for this combination is CK

MIXING COCAINE AND OTHER DRUGS

Alcohol and cocaine

Causes toxic reactions that increase heart attack and stroke risks

Heroin and cocaine

Raises your risk of overdose, injecting damage and drug dependence

Benzos and cocaine

Increases your risk of overdose, confusion and risky decision making

Ketamine and cocaine

Triggers psychotic episodes, heart attacks, strokes, and bladder problems

Cocaine compulsion and psychological dependence

The compulsive nature of cocaine is related to the powerful effect it has on the brain's reward system. People can feel an overwhelming, and sometime unexpected urge to use, even after a significant period of abstinence. Cravings are often triggered by sights, sounds, smells or using other drugs and alcohol.

Daily use can become an integral part of people's lives and they can find functioning (physically, psychologically and socially) without it very difficult. Although no physical dependence occurs, even with repeated use, the psychological grip is very real and should not be underestimated. It should be noted that the symptoms of anxiety, which are often present after a binge or on cessation of use, will very much feel to the person like physical withdrawals.

Cocaine withdrawal (sometimes called a 'come down') from heavy cocaine use can be very uncomfortable. Cocaine withdrawal is however rarely life threatening.

A commonly cited study into cocaine withdrawal was undertaken by Gawin and Kleber in 1986 and is still relevant today. Using data collected from 30 cocaine-dependent outpatients, the researchers reported three distinct phases of the withdrawal process: 'crash, withdrawal and extinction':

We have developed this model further to include high-risk times for other substance use which may increase the likelihood of lapse or relapse.

Phase one, 'the crash' develops rapidly following abrupt cessation of heavy cocaine use and is characterised by acute dysphoria, irritability and anxiety, increased desire for sleep, exhaustion, increased appetite, decreased craving to use. **Other substances may be taken to help ease the comedown.**

Phase two, 'withdrawal' is characterised by increasing craving to use, poor concentration, some irritability and lethargy, which persists for up to 10 weeks. **Other substances may be taken to cope with negative feelings or deal with cravings.**

Phase three, 'extinction or eradication' comprises fewer intermittent cravings to use in the context of external cues. **These cravings may be at a stage where they are easier to deal with without the use of other substances.**

COCAINE WITHDRAWALS



Stages of cocaine withdrawal and recovery

Some people choose to take other drugs, such as alcohol, sleeping tablets, opiates or benzodiazepines, to help ease the crash and withdrawal described in the first 10 weeks following cessation of cocaine use. This can lead to additional problems including dependence on different drugs, drug related harms and the risk of overdose.

It may be the case that a person will continue to use cocaine even though negative consequences are evident. This is a good indicator of drug dependence, albeit psychological. Binge use of cocaine is common and can quickly lead to a range of physical and mental health problems. It may be however, that social and financial problems appear before the physical or mental health consequences of cocaine use.

Cocaine harms and harm reduction

Regardless of how cocaine is administered, there are a wide range of potential significant harms.

Cocaine Overdose Stroke Heart attack Toxicity (COSHT)

Cocaine was implicated in almost half of the drug related deaths in 2023. At toxicology, cocaine is often found to be present with other drugs. These other drugs are usually central nervous system depressant (CNSD) drugs such as heroin or methadone. For this reason, naloxone should always be administered when a more typical CNSD overdose presents where cocaine has also been taken.

The signs and symptoms of a cocaine overdose can be different from those we expect to see in a typical CNSD overdose because cocaine is a stimulant drug.

A cocaine overdose occurs when excessive cocaine intake overwhelms the body causing a life-threatening state. This can result in toxicity, stroke and heart attack.

The mechanism of COSHT involves a complex interaction of pharmacological effects leading to multi-organ dysfunction. This should always be treated as a medical emergency.

- **Dopamine and norepinephrine reuptake blockade**- This prolongs their duration of action in the synaptic clefts and results in excessive sympathetic nervous system activation (fight-or-flight response), causing tachycardia (fast heart rate), hypertension (high blood pressure), hyperthermia (high body temperature), and agitation.
- **Serotonin reuptake blockade**- Whilst less prominent an impact than dopamine or norepinephrine, serotonin reuptake inhibition may contribute to CNS excitation and potential neurotoxicity (damage to the brain).
- **Sodium Channel Blockade**- Cocaine acts as a local anaesthetic by blocking sodium channels in cardiac tissue. This impairs electrical conduction leading to arrhythmias (irregular heart rhythm) or depressed cardiac function.
- **Vasoconstriction (the narrowing of blood vessels) and ischemia (sub optimal flow of blood)**- This coronary vasoconstriction can precipitate myocardial infarction (heart attack), while cerebral (brain) vasoconstriction increases the risk of stroke.
- **Platelet Activation**- Cocaine enhances platelet aggregation (the process where tiny blood cells stick together to form a clot or blockage). This increases the likelihood of obstructing blood flow meaning heart attack or stroke become a risk.
- **Central Nervous System (CNS) Effects**- Overstimulation of the CNS leads to seizures, hyperthermia and agitation. Hyperthermia also exacerbates metabolic demand and can cause muscle breakdown leading to acute kidney damage.

These mechanisms lead to the life-threatening conditions of COSHT such as arrhythmias, heart attack, stroke, seizures, kidney failure, and multi-organ collapse. Rapid medical attention will increase the likelihood of a positive outcome.



Effects on the body:

- Over stimulates the brain and heart
- Raises blood pressure, heart rate and temperature
- Can cause heart attacks, stroke or seizures

Warning signs:

- Chest pain, pounding heart, over heating, collapse
- Stroke signs like drooping face or weak limbs
- Confusion, hallucinations, extreme agitation

What to do:

- Call emergency services immediately on 999
- Ask someone to stay with you
- Don't use any more drugs or alcohol

Signs and Symptoms of COSHT

- Stroke signs (drooping of the face or loss of movement in the limbs)
- Heart attack signs (pain in chest, arms, jaw, neck)
- High blood pressure (headache, dizziness)
- Increased heart rate (pounding chest, palpitations)
- Severe headache
- Blurring or loss of vision
- Severe agitation, restlessness, confusion, excited delirium*
- Hyperthermia (sweating not caused by heat or exercise)
- Abnormal reflexes/movement
- Seizure
- Diarrhoea, vomiting, abdominal pain

If COSHT is suspected the person should call emergency services as soon as possible and follow the call handler's instructions.

*Cocaine induced excited delirium. People displaying symptoms such as aggression, hyperactivity, extreme paranoia, hyperthermia, incoherent shouting or screaming and unusual strength are often at risk for sudden death.

Clinical Outcomes

Management within a hospital setting will focus on stabilizing vital signs and may include treatments such as cooling for hyperthermia, benzodiazepines for agitation/seizures and medicating other specific complications.

Reducing the Risk of COSHT

- Taking less cocaine (dose)
- Lengthen the time between doses
- Limit the duration of the sitting
- Not consuming other drugs, alcohol or tobacco within the same sitting
- Accessing heart health checks as a means of identifying cardiovascular problems at the earliest possible stage

Cocaine and sexual risks

Like many other drugs, cocaine can lower inhibitions and make a person more likely to engage in sexual activity. This, often impulsive sex, may reduce the likelihood of barrier contraception use or other safe sex practices.

Rougher sex, such as anal sex, may seem more appealing whilst prolonged sex is common due to the inability to climax. This may lead to the tearing of membranes causing bleeding and increases the potential to transmit blood borne viruses (BBV).

Men may find it difficult to get and maintain an erection, leading to a frustrated state.

Overall, there are considerable risk factors associated with cocaine use and sex. Unplanned pregnancies, sexually transmitted infections (STIs), BBV's and sexual assault are some of the potential negative consequences of having sex under the influence of cocaine.

Longer term cocaine use can change a woman's menstrual cycle and stop ovulation, whilst damage to the fallopian tubes from STIs can result in infertility.

Cocaine use during pregnancy can cause seizures, migraines, premature birth and in some cases, the placenta can detach from the uterine wall (placental abruption). Stillbirths and miscarriages occur at higher rates in women who use cocaine during pregnancy.

COCAINE AND YOUR SEXUAL HEALTH

Using cocaine can affect your judgement and make it harder to give or recognise clear sexual consent.



Risky decisions - you are more likely to have unsafe sex e.g. no condom	Rougher or longer sex can cause cuts and bleeding and increase risk of infections	Cocaine can cause erection issues, fertility problems and disrupt periods
People using cocaine are more likely to get an STI or become pregnant	Some people trade sex for cocaine, increasing chances of assault or abuse	Using when pregnant can cause miscarriage, early labour and harm the baby



Cocaine, Malnutrition and the Immune System

People taking cocaine, especially during binges, often neglect basic self-care. Regular meals, hydration and rest can become secondary to the compulsion to use. Cocaine suppresses appetite, increases energy expenditure and disturbs sleep patterns, all of which can contribute to significant nutritional deficits over time. These effects are especially concerning for people who use heavily or regularly and for those who are already in vulnerable physical health due to homelessness, stress, or poly-substance use.

The combination of cocaine and alcohol is particularly harmful to the body. While alcohol contains calories, it lacks the nutrients needed to maintain health. Alcohol is also a powerful diuretic, meaning it increases fluid loss through urination and contributes directly to dehydration. When people are drinking and using stimulants at the same time, they often do not feel thirsty or may forget to drink water entirely. This significantly increases the risk of becoming dehydrated, especially during all night sessions or binges. The body becomes depleted of not only water, but also vital electrolytes like potassium and sodium, which are essential for heart, kidney, and muscle function.

Cocaine use places considerable stress on the body. Elevated heart rate, increased body temperature, and prolonged periods without rest deplete energy reserves and further dehydrate the system.

Binge use may involve hours or days without food or water, followed by vomiting, diarrhoea, or exhaustion during the crash. These cycles disrupt the body's ability to absorb nutrients and repair itself, weakening both physical resilience and immune defence. Over time, this increases the risk of infection, slows wound healing, and worsens the impact of other health conditions.

Malnutrition also affects the brain. A lack of essential nutrients such as B vitamins, magnesium, and zinc can worsen mood instability, anxiety and cognitive issues. People experiencing poor nutrition may find it harder to cope with cravings, recover after binges, or regulate their emotions. When combined with sleep deprivation and alcohol related toxicity, these effects can intensify mental health symptoms and increase the likelihood of continued or riskier drug use.

Supporting Better Nutrition and Self Care

Supporting people who use cocaine to maintain even small, manageable routines around eating and drinking can make a real difference. This does not need to mean large meals. Snacks with protein, fruit, or slow release carbohydrates are often more achievable, especially during or after a binge. Encouraging people to keep a bottle of water nearby and take regular sips can reduce dehydration, headaches, and overheating, which are common features of stimulant use.

Before a cocaine session, particularly if use is planned in advance, there is an opportunity to frontload on nutrients and prepare for hydration. Eating a balanced meal beforehand can help reduce the strain on the body, especially if food is unlikely to be consumed during the session itself. Isotonic drinks (such as sports rehydration fluids) are particularly helpful, as they replace both fluids and essential electrolytes that are quickly lost during stimulant use.

This planning stage can also be used to stock up on easy, good quality nutrition. Having ready meals, fruit, protein rich snacks, and hydrating drinks in the fridge or cupboards may make it easier to eat and recover during the comedown phase, when energy is low and the person may not feel like shopping or preparing food. Even small steps like this can improve recovery and reduce longer-term health impacts.

Services and supported accommodations may wish to consider the role that nutrition and hydration play in reducing harm. When there is awareness that cocaine is being used, particularly in binge patterns, staff should reflect on the quality, timing, and availability of meals and fluids. Even small adjustments, such as extending food access hours or providing out of hours nutritious may support recovery and reduce health risks for those using cocaine.

COCAINE AND NUTRITION



Mouth ulcers



Nausea



Weight loss



Dehydration



Malnutrition

Using cocaine causes you to lose weight

Malnutrition can affect your immune system, bones, muscles, sleep and mental health

Risks associated with methods of administration

Snort risks

Septum damage is one of the most significant long-term effects of cocaine snorting. A septal perforation, or "hole in the septum", is a condition commonly caused from this method of administration.

The nose has a fragile blood supply which is reduced by cocaine snorting. This process is called vasoconstriction (narrowing of blood vessels). When the blood vessels constrict, the blood supply is compromised, delivering less oxygen to the tissues of the septum. With low oxygen, the septum lining begins to die. Once the lining is damaged, it can no longer support the cartilage underneath and the cartilage dies. This is called a septal perforation. Once the septum is perforated, the nose can collapse because the septum is the structural support of the nose. **Once a septal perforation is present it will never heal on its own.**

It is much easier to deal with septum damage if it is identified early. Medical advice should be sought at the earliest opportunity if any of the following symptoms occur: constant runny nose, discharge from the nose or frequent bleeding

Typical lines of cocaine ready for snorting



COCAINE AND YOUR NOSE



Snorting cocaine can damage your nose



It stops blood flowing, damages soft tissues and wears away the protective lining



Causing nose bleeds and even a hole between your nostrils

Transmission of infections including blood borne viruses

Straws, tubes or banknotes inserted into the nose can come into contact with Hepatitis C or Hepatitis B infected blood. This can be transmitted to someone else sharing them. The risk is probably lower than previously thought, however should still be taken seriously. Sharing snorting items will also have a risk of transmitting other infections or viruses, including Covid.

Snort harm reduction

- Not all white powder is cocaine and therefore a smaller line should be taken as a test dose
- Dosage should be timed, leaving at least 30-45 min between lines
- Lines should be alternated between both nostrils, unless one is damaged or bleeding
- The surface the line is snorted from should be as clean and sterile as possible. A wipe or spray with an antibacterial agent will help this
- As cocaine varies in purity so too should the line size (dose). The purer the cocaine the smaller the line/dose should be
- The powder should be crushed or chopped as finely as possible to save blocking the nose
- A straw, tube or plastic type banknote which is long enough to reach high up into the nose should be chosen. This should be unused and not shared with others
- Old type paper bank notes are grubby by nature and difficult to clean and so these should be avoided
- If snorting with others the tube should be taken and kept on the person (in pocket, bag etc.) until next use to reduce accidental sharing
- Blowing the nose when a build-up of mucus is felt will help clear residue
- Dousing (sniffing up water from the palm of the hand then blowing the nose and repeating) after the session is complete will help clear any residue

Smoke risks

Prolonged crack/freebase smoking is likely to result in some form of lung problem. Repeated inhalations can cause or exacerbate lung or respiratory conditions such as asthma, shortness of breath or chronic obstructive pulmonary disease (COPD). Smoking crack cocaine may also cause pneumonia.

Crack lung is a clinical condition where alveolar hemorrhage (bleeding) occurs acutely within 48 hours of smoking crack or freebase cocaine.

Crack lung has also become a term used to describe any acute pulmonary injury related to smoking crack cocaine. Some people may experience such severe damage to the lungs that they spit up blood, whilst serious cases may result in respiratory failure.

The chemicals used during the preparation process can be inhaled directly into the lungs causing harm. This is particularly relevant when smoking freebase rocks where the ammonia has not been properly rinsed before smoking.

Symptoms such as chest pain, extreme coughing fits and difficulty breathing are all indications that significant harm is occurring.

COCAINE AND YOUR LUNGS



Burns



Infections



Bleeding



Scarring



Toxic damage

Smoking crack or freebase damages your lungs

It affects your breathing and can make lung problems like asthma worse

Pipe risks

Ideally people would smoke their crack/freebase from a purpose made pipe or glass/Pyrex tube with suitable gauze. If these are not easily accessible people will make their own from easy to find household items. Drinks cans, plastic bottles, glass miniature bottles and inhalers are all commonly used. A bed of cigarette ash can be used to better melt the rock without clogging the holes, however the inhalation of this burning ash can create further risks. The risk associated with each type of pipe varies greatly. If any of these pipes are shared then there is a potential for a wide range of viruses and infections to be transmitted including BBVs, Tuberculosis and Covid etc.

Drinks cans- Holes are made and the rock is often placed on a bed of cigarette ash. Toxic fumes from the burning paint or plastic coating inside the can may be inhaled along with the drug itself. There is a risk of virus transmission and other infections if shared with others.

Plastic bottles, inhalers and other plastic items- Fumes from burning plastic can be inhaled along with the drug. Some of these pipes place the rock and flame very close to the face resulting in burns to the head and neck. There is a risk of virus transmission and other infections if shared with others.

Glass miniature bottles- The bottom is often taken off of this small bottle leaving sharp edges which may cause cuts. There is a risk of virus transmission and other infections if shared with others.

Gauze- It is common for loose filters to be made from stainless steel scouring pads. These are chosen as a bed for the rock to sit on allowing better melting and inhalation. However, these can disintegrate and break off during heating, meaning small particles can be inhaled causing significant damage to the mouth, throat and lungs.

A crack pipe provided by harm reduction services (not UK)



A makeshift crack pipe made from a salbutamol inhaler



Recycling cocaine from pipes

When a pipe has been used numerous times cocaine residue will gather inside. In order to free this residue for further use, a chemical such as acetone or nail varnish remover can be used to dissolve the cocaine. The liquid can then be poured on to a mirror and left to dry. The cocaine recovered can be scraped off of the mirror and re-used. Inhaling these chemicals before they have fully evaporated can cause lung damage.

Smoke harm reduction

Personally made freebase cocaine is less likely to have impurities when compared to street bought crack. Freebase rocks should be rinsed properly to wash off any ammonia residue. Placing rocks on a damp tea bag and carefully rinsing with cold water before allowing to dry prior to smoking will help to remove ammonia.

Purpose made toughened glass or steel pipes are likely to be the safest choice. The pipe should be long enough that it is far enough away from the face to stop any burns from the flame or debris.

Pipes should not be shared with any other person to reduce the risk of virus transmission and infection.

Proper pipe gauze should be used in place of metal scouring pad type material. This should be carefully inserted into the pipe or tube and changed frequently.

A lighter which has enough reach to melt the rock without burning the fingers should be used.

Use of a blow torch poses a significant fire risk if the person collapses as it will not shut off automatically. In this scenario people should be advised not to use alone.

Holding the crack or freebase smoke in the lungs for long periods can damage the lung tissue without increasing effects of cocaine.

Smoking tobacco along with cocaine increases strain on the heart and cardiovascular system.

Any burns, blisters, sores or chaps on the lips should be treated properly with appropriate ointment or cream.

Good hydration through drinking lots of fresh water may stop the lips becoming dry and chapped.

Injecting related harm

Injecting cocaine is likely to be the most harmful method of administration.

Blood Borne Virus (BBV) risks

Sharing needles and syringes carries the greatest risk of transmitting BBVs (Hepatitis B, Hepatitis C and HIV). Other injecting paraphernalia such as spoons, filters and water also carry a risk if shared, albeit it to a lesser degree. The indirect sharing of injecting paraphernalia is common, particularly if batches of drugs are prepared using previously used equipment. Drawing the drug solution from a contaminated spoon, filter or water has both bacterial and BBV infection risks. It is also common for all the drugs to be drawn into one syringe and then divided equally by back-loading or frontloading into people's syringes, which again creates infection transmission risks.

Injecting related complications and injury

The effect of cocaine is short lived and so the compulsion to inject frequently occurs. This can lead to a "binge injecting session". This frequent injecting can cause rapid deterioration of the veins, leading to vein collapse and circulatory problems. The anaesthetic effects of cocaine can make it difficult to 'feel' the injection properly, leading to missed hits and site damage.

Subcutaneous or intramuscular injecting under the skin (sometimes called skin or muscle popping) can cause significant damage to the skin, tissue and muscle. The muscle breakdown can cause toxins in the blood which cause damage to the kidneys. This can result in a rare condition called rhabdomyolysis

It may be difficult for people to estimate how many times they are likely to inject in any given session. This can result in the person not collecting enough injecting equipment and so having to reuse. Even the reuse of someone's own (not shared) injecting equipment can result in bacterial infection, vein and injection site damage.

It is possible that the frequency of injection, often across multiple geographic locations, is a driving factor in poor general hygiene and unsterile injection practices. This has been shown to cause a number of bacterial infections such as *Staphylococcus aureus* and Group A Streptococci. Infections can result in serious life threatening conditions such as sepsis or necrotizing fasciitis.

INJECTING COCAINE



Injecting cocaine can damage your veins and soft tissues



Causing skin infections and a risk of hepatitis and HIV



It is the most dangerous way to take cocaine

An outdoor/away from home public injecting site close to Glasgow City Centre.




Preparing cocaine for injection

Each form of cocaine is prepared for injection slightly differently.

Powder cocaine is easily water soluble without the need for an acidifier or heat. Freebase and crack do need an acidifier added to facilitate the breakdown into a soluble state. **Crack and freebase should not be heated** as this will form an oil which may block the needle or cause damage if injected.

Snowballing (heroin and powder cocaine together in the one syringe) would be prepared on the spoon by cooking up the heroin as normal (using an acidifier and heat). When the solution is cooling, powder cocaine is added.

Snowballing with heroin and crack or freebase is a little trickier. The crack or freebase should be crushed as fine as possible before adding cold water and acidifier. When dissolved, heroin should be added and the solution heated. More acidifier can be added in very small amounts if needed.




Cocaine injecting harm reduction

- Injecting equipment such as needles, syringes, filters and spoons should not be shared with others
- All injecting equipment such as needles, syringes, spoons, filters and water should be new and unused
- All injecting equipment should be considered single use and disposed of in a suitable sharps bin immediately after use
- The drugs should be prepared and injected in as sterile an environment as possible. Ideally this environment should also be warm and well-lit with other trusted people there to help in an emergency
- Hands and target injecting sites should be washed well with warm water and soap. If this is not possible the best available method of cleaning should be used
- Techniques for raising veins (tourniquet, gentle exercise, warm water) should be used to keep lower risk sites accessible for longer
- The size of needle should be carefully chosen in relation to the intended injection site. The smallest possible needle should be selected. Deep vein injecting will require a longer more robust needle
- If batches must be prepared with others, all equipment that every person uses should be new and unused
- If preparing powder cocaine for injection then no acidifier or heat is required. Acidifiers cause vein damage and so should only be used if essential
- Proper rotation of injection sites is advised. Ideally 5 or 6 low risk veins on the arm should be identified and kept clean in between use
- Injecting intramuscularly or subcutaneously should be avoided as this carries a significant risk of skin, tissue and muscle damage

Needle and Syringe Guide

Colour and description	Gauge	Length	Suitable for	Drugs normally injected with this size	Available in Single Use Packs
Fixed needle and syringe	29g	1 Inch	Superficial veins, arms, hands, feet	Heroin, cocaine, steroids, powder hormones	Yes (black writing)
Blue	23g	1 ¼ Inch	Femoral vein, intramuscular	Heroin, cocaine, steroids	Yes (blue writing)
Blue	23g	1 Inch	Femoral vein, intramuscular	Heroin, cocaine, steroids	No
Orange	25g	1 Inch	Femoral vein, other partially deeper veins	Heroin, cocaine	Yes (orange writing)
Orange	25g	5/8 Inch	partially deeper veins	Heroin, cocaine	No

Example of single use pack: black containing a fixed 29-gauge needle and syringe, spoon/cooker, filter, vitamin C and pre injection cleansing swab



Physical health monitoring and treatment interventions for individuals who are prescribed OST and are using cocaine

An update of the physical health history and regular review of all prescribed medication is particularly important for individuals using cocaine as there is greater risk of strain on the heart from medications which prolong the QT interval. There is also an increased risk with sedative medications, both prescribed and non-prescribed, that cocaine may mask the depressant impact on the respiratory system and so increase risk of overdose.

There is evidence that participation in treatment and commencement of opioid substitution therapy (OST) can be associated with reduced cocaine use in individuals with opiate dependence also using cocaine. It would therefore not be advisable to delay OST in an attempt to stabilise cocaine use first.

An ECG can give a helpful indication of strain on the heart and any prolongation of the QT interval. Where possible this is helpful at initiation of OST or during changes in dose. If this is not possible a balanced judgement is required as holding up vital treatment to await an ECG first may increase rather than reduce risk to health.

There is reduced risk of QT interval prolongation (adding to heart strain) and reduced respiratory depression risk with buprenorphine compared to methadone. Although there is no specific recommendation in OST choice for individuals with opiate dependence who are using cocaine, this should be taken into consideration along with other individual factors.

Individuals who start to use cocaine during OST may require an update in their risk and Red/Amber/Green (RAG) assessment and more regular support from their care manager. Consideration should be given to instalment dispensing of OST.

Cocaine use should not prevent an increase in OST if this is indicated. Stabilising on an optimal dose of OST may not only make it easier for the individual to reduce and stop using non-prescribed opiates but also other drugs such as cocaine.

COCAINE AND YOUR HEART

Cocaine makes your heart beat faster and raises your blood pressure, while letting less oxygen reach your heart. It messes with your heart's rhythm, makes your blood clot, and can harm your heart muscle.

- ➡ **Heart attacks**
- ➡ **Heart rhythm problems**
- ➡ **Heart failure**
- ➡ **Ripped blood vessels**
- ➡ **Narrowed blood vessels**
- ➡ **Heart valve infections**



Mental health effects and treatment interventions

Crisis intervention

A crisis is an acute, time-limited episode experienced as overwhelming emotional reactions to an event. What is a crisis for one person may not be so for another. What becomes a crisis may not have been a crisis before or would not be a crisis in a different setting. Crisis has been described as a system out of balance. Crises occur when balance cannot be regained, even though a person is trying very hard to correct the imbalance.

Symptoms of drug related mental health crisis

It can be difficult to differentiate between the symptoms of mental illness and temporary drug-affected behaviour. Therefore, it is often not possible to conduct a mental health assessment until the effects of the drugs have fully worn off. However, it may be necessary to provide a crisis assessment in order to ascertain if an immediate mental health intervention is required due to temporary incapacity or where there is a risk to the person or others.

Common symptoms of drug related mental health crisis

- Extreme anxiety/paranoia
- Hallucinations- auditory (hearing things), visual (seeing things), and tactile (bodily sensations such as feeling things crawling on them)
- Persecutory beliefs (believing someone is trying to harm them)
- Delusional thoughts (believing they are someone else or they have done something they have not)
- Self-harm
- Suicidal ideation/acts
- Loss of grip on reality
- Whilst some people report being concerned about aggression, this is usually a fear response and is unlikely to occur if the person is appropriately supported

In the majority of cases symptoms resolve as substance(s) wear off and where the person is able to sleep.

Sometimes with severe drug toxicity, delusional behaviours and extreme agitation can occur in the presence of other serious physical health symptoms such as extreme overheating and extremely fast heart rate. Delusional behaviours may manifest as bizarre behaviours e.g. getting undressed in public or aggression and extreme fear; this condition is referred to as acute behavioural disturbance or excited delirium and is a medical emergency. In the case of stimulants, excited delirium is caused by a condition called serotonin syndrome and can be life threatening if left untreated or if the person is placed in restraints/held down with force for prolonged periods.

Therefore, if a stimulant related psychosis is suspected e.g. person presents with excited delirium symptoms, it is important to rule out a stimulant overdose. Signs to look out for are extreme overheating (being hot to touch and/or profusely sweating), muscle rigidity, over responsive reflexes or involuntary muscle spasms e.g. jerking. Other signs might be seemingly insensitive to pain, displaying a high level of physical activity /energy without tiring and appearing to have excessive strength (this can be due to fear if struggling against being contained or against use of restraint). If such physical symptoms are identified, urgent medical care should be sought by phoning **999**.

Where there are no signs of the physical symptoms connected to excited delirium, cases can be managed as any acute mental health crisis. Focus would be on providing a safe environment, managing agitation and offering reassurance and de-escalation in the first instance.

COCAINE AND YOUR BRAIN

Cocaine changes the chemical levels in the reward part of your brain

Short term dopamine high:
Extra dopamine produces short lived feelings of euphoria, extra energy, social confidence and well-being

Come down crash:
After using, low chemical levels cause extreme tiredness, anxiety, irritability and low mood

Mental health risks:
Long term use can cause serious mental health problems like paranoia and psychosis, which may need psychiatric treatment

Responding to a drug induced mental health crisis

De-escalation

For people experiencing an acute drug related mental health crisis, simple de-escalation techniques are often highly effective. These include:

Techniques to establish safety and trust:

- Speak slowly and confidently with a gentle, calm tone of voice
- Give your name and explain your role and purpose if you do not know the person
- Use non-threatening body language
- Avoid touching, shouting or sudden movement
- Stay calm and provide any necessary support slowly and gently

Offer clear and supportive communication:

- Use clear language
- Explain any actions you need to take beforehand
- Avoid intense questioning
- Ask how you can best help the person right now
- Paraphrase their concerns
- Do not challenge psychotic thinking or collude in delusions
- Do not argue or threaten
- Where someone is presenting as mentally confused avoid sarcasm, laughing or humour that may be misunderstood
- Comply with reasonable requests e.g. making a phone call

Establish physical safety:

- Think about environment. Is there somewhere quieter or where a person might feel more comfortable? Be aware of risk management when staying in or changing an environment
- Reduce unhelpful distractions/audiences (ask others to leave)
- Consider if having a friend (s) to stay with someone is helpful/unhelpful
- Avoid restricting the person's movement unless unsafe

Verbal reassurance and support:

Talking through or down, depending on an individual's drug experience and levels of agitation. It may be more appropriate to simply empathetically listen to concerns and offer reassurance where required. Where someone is very agitated or less experienced, they may prefer to be talked down where you are more directive in your support. Some useful strategies for this are:

- Normalising their experience e.g. "cocaine can cause some people to feel anxious or paranoid, you are safe here and the effects will start to wear off soon"
- Distraction and re-focusing techniques e.g. engaging in a conversation about something else or asking them to focus on something in the room
- Grounding techniques e.g. deep belly breathing, counting (an example techniques is described below)

Five things grounding technique

- Notice five things that you can see. Look around you and bring your attention to five things that you can see. Pick something that you do not normally notice, like a shadow or a small crack in the concrete.
- Notice four things that you can feel. Bring awareness to four things that you are currently feeling, like the texture of your trousers, the feeling of the breeze on your skin, or the smooth surface of a table you are resting your hands on.
- Notice three things you can hear. Take a moment to listen, and note three things that you hear in the background. This can be the chirp of a bird, the hum of the refrigerator, or the faint sounds of traffic from a nearby road.
- Notice two things you can smell. Bring your awareness to smells that you usually filter out, whether they are pleasant or unpleasant. Perhaps the breeze is carrying a scent of trees if you are outside, or the smell of a fast food restaurant across the street.
- Notice one thing you can taste. Focus on one thing that you can taste right now, in this moment. You can take a sip of a drink, chew a piece of chewing gum, eat something, or think of something you like the taste of.

Next steps

Once a level of safety has been established and de-escalation strategies have helped the acute phase of the crisis to pass, it is important to look at what other supports might be beneficial. The person may benefit from support to develop self-management techniques should future crises occur.

It is common for people, particularly where they have underlying mental health issues, to experience repeated crises. These experiences can provide insights into resources and coping skills that have been effective or have yet to be tried. Discussions can help formulate action plans and form self-management techniques.

Some people may require more specialist help as part of follow-up. This may include referral to community mental health or psychological therapies teams (see Treatment section).

The acute effects of cocaine can include a sense of inflated confidence, sexual arousal, reduced need for sleep, reduced appetite and increased agitation and paranoia. Taking cocaine when already feeling anxious can heighten the sense of anxiety. Deprived sleep can exacerbate existing mental health difficulties as well as cause new mental health problems.

Mental Health Assessment

As described above, mental health symptoms usually resolve over hours or days after stopping use of cocaine, but they may be more persistent.

Chronic use or heavy binges can lead to the development of paranoid ideation associated with anxiety, panic attacks and feeling agitated. Cocaine use can also lead to depression and suicidal thoughts. Initial anxiety and paranoia may sometimes progress to a psychotic disorder with paranoid delusions and hallucinations (which may be auditory or tactile "cocaine bugs").

In some cases, individuals can present only with delusional parasitosis, Ekbom Syndrome, which is the unshakable belief that the skin is infested by parasites in the absence of other mental health symptoms. Characteristic is the self-damage of skin by individuals affected in an attempt to remove parasites. Presentation to primary care and A&E for treatment rather than to mental health or addiction services is common. To prove infestation, containers with specimens or photos ('matchbox sign') may be brought to consultations. Psychological distress secondary to delusions can lead to depressed mood and may require further assessment and/or treatment.

When working with people using stimulants, it can be helpful to screen for mental health problems. It is important to enquire about sleep patterns, anxiety, fears, delusional beliefs or hallucinations. People should be asked about thoughts of self-harm and suicide. Any concerns should be appropriately escalated. A particular focus on sleep hygiene, anxiety management techniques and building a safe environment to recover from acute effects of cocaine can be helpful.

Persistent mental health symptoms beyond the acute effects of cocaine should be treated as they arise. Awareness of interaction risks when prescribing medication to people who use cocaine should be considered. A specific consideration is that cocaine can mask the depressant effect of sedative medication. People should be aware of risks of respiratory depression and need for additional caution when using sedatives alongside cocaine. Selective serotonin re-uptake inhibitors (SSRIs) and stimulants have been known to produce toxic reactions, although not commonly. The Medicines and Healthcare Products Regulatory Agency (MHRA) has reminded prescribers to note the potential increased risk of bleeding when citalopram is prescribed to patients who are taking cocaine.

Cocaine use can lead to acute psychiatric presentations which require emergency assessment and possible admission. This would be applicable if a person presents with symptoms of paranoia, hallucinations and in a state of agitation after the acute effects of cocaine have worn off, or if a person presents with acute distress leading to suicidal ideation.

Some helpful contacts when further emergency assessment or admission may be required or to form part of a safety plan for a patient:

Breathing Space: ☎ **0800 83 85 87** (a free confidential service which offers telephone support to people experiencing low mood, depression and anxiety).

NHS 24 Mental Health Hub can be accessed by calling ☎ **111**. They are available 24 hours a day for patients in mental health crisis and hub staff will then support access to the OOHs CPN Service if specialist mental health assessment is required. This provides people with one point of access and no call charges. They can also be supported to access the most appropriate response to meet their needs at the time they need it.

In addition, there is direct access to the Out of Hours CPN service via the Mental Health Assessment Units based at Leverndale and Stobhill Hospitals, which can be accessed through the NHS GG&C switchboard or by calling ☎ **0141 211 3600**. This number is for profession-to-profession referrals only and not for general public use.

Where there is uncertainty around roles of addiction workers and Adult Mental Health colleagues it may be helpful to refer to the following document which can be found on the MyPsych website & app: [**Adult Mental Health & Addictions Service Shared Guidance and Specification for Interface Working March 2021**](#).

Psycho-social Interventions

The three common parts of effective psychosocial interventions for substance use such as crack cocaine and heroin / OST are: **enhancing motivation, developing control over impulsive behaviour and developing a rewarding lifestyle***.

Stage 1: Motivational Enhancement

Motivational enhancement is an essential component of any intervention of this kind. Different strategies will be utilised depending on peoples' level of engagement. In the earlier stages of engagement, the first and second strategies (below) may be more useful. As the engagement deepens and strengthens, the later strategies will become more useful. However, people's engagement will fluctuate and it is important to ensure the correct strategies are being utilised at the right time. The main motivational enhancement strategies include:

1. Establishing a supportive, caring therapeutic relationship. Establishing a relationship and developing communication skills is important to enable a conversation about drug use. Effective therapeutic relationship may be strengthened by helping the individual to identify practical needs, problems and priorities. Crisis intervention and stabilisation of acute symptoms may help to strengthen the relationship.
2. Finding topics that motivate interest (motivational hooks). Find out what is important to the individual; what do they want to change in their lives. Explore their relationships and goals to find topics that motivate interest.
3. Enabling conversations which explore and discuss substance use. Why are they using? What do they like about the drug? What role does it have in their life e.g. do they think it helps them to cope with mental health difficulties. Try to remain neutral. People will be more likely to listen to information on the adverse effects of the drug, if they feel they have been listened to. Before providing this information, check their existing knowledge i.e. individuals are often aware of the risks of drug use.
4. Explore the pros and cons of behaviour change. Firstly, explore the pros and cons of their current drug use, then move on to the cons and pros of making the change.
5. Strategies for supporting positive change. Look out for statements of concern or intent to change. Ask readiness to change questions. Rate the importance of each pro and con. Identify and question the positive beliefs around drug use.

Stage 2: Developing control over impulsive behaviour

Developing control over impulsive behaviour begins with investigating the behaviour itself (e.g. why someone smokes crack cocaine), in addition to what was happening before the drug use, and what the consequences of the drug use were. This process is known as Functional Analysis and it can help an individual to realise that when a trigger occurs, they can use a non-drug using behaviour, resulting in a different set of positive long-term consequences. This is one of the core principles of controlling impulsive behaviour, which involves **identifying and managing triggers, coping with cravings, analysing seemingly irrelevant decisions and managing high risk situations**.

Identifying and managing triggers

Individuals may not be aware of their triggers, or even that they have them. Identifying triggers enhances control in that it is the first step in learning how to manage triggers. A detailed drug diary may help to identify their triggers (as well as understanding the links between triggers, drug use and consequences; and potentially to develop a better sense of control over their drug use). The Discovering Triggers worksheet can also help, as can asking the following questions:

- Before the individual used the drug, what were they thinking, feeling, doing; where were they, who were they with?
- What was the perceived coping mechanism e.g. did they believe that it improved their feelings of wellbeing or alleviated boredom?
- What were the positive and negative short and long-term consequences?

In terms of managing triggers, the individual should try to remove all of the possible environmental triggers. This will be helped by the Self-Management Plan. For those triggers that cannot be removed, develop a craving plan, using the Coping with Cravings and Urges worksheet. At some point in the future, some environmental triggers may be able to return, as the coping strategies strengthen. It is also important to identify thoughts and feelings which are triggers and develop a coping plan, using the Coping with Thoughts worksheet.

Coping with cravings

The following activities may help individuals to manage their cravings:

1. Distraction involves any activity that diverts their attention from the drug craving and may involve mental, physical or soothing activities. Examples include participating in sports; rediscovering interests such as art, music, film; gaming, doing crosswords, Sudoku, other puzzles, reading, TV etc.
2. In a safe setting, focus on the craving and find ways to cope with the feeling such as urge surfing, mindfulness or recovery meetings.

Again, a drug diary will be helpful in managing cravings.

Seemingly irrelevant decisions and high risk situations

Seemingly irrelevant decisions occur when a series of decisions are made, that increase the risk of lapse or relapse. For example, going to the local shop and meeting an old friend may put the individual at risk of a lapse or relapse. The Seemingly Irrelevant Decisions Worksheet will help individuals gain a broader understanding of these. The individual should be supported in developing an awareness of high risk situations in terms of people, places, thoughts and feelings.

Stage 3: Developing a rewarding lifestyle

When an individual has mastered many of the skills above and has developed a strong sense of control over their substance use, it is important to support them in developing a more rewarding lifestyle, as this will help strengthen their recovery. Supporting an individual to develop new or existing skills, may help improve motivation, encourage greater stability and ensure they have the commitment and energy to develop holistically. It is important to help someone develop new experiences and explore ways in which they pursue new interests. The exercises and worksheets highlighted below can help to structure the conversation and develop ideas:

1. Set priorities using the Happiness Scale and identify person-centred activities using the Recreational Survey.
2. Set goals using the Goal Setting Worksheet.
3. Monitor progress using the Highlighting Progress Graph and Calendar.

**This guidance has been taken from the NHS Education for Scotland (NES) Core Behavioural and CBT Skills for Relapse Prevention and Recovery Management Course (Laura Freeman, NES, 2011). In order to follow and implement this guidance effectively, it is recommended that workers should have attended Core Skills training. The worksheets mentioned in this guidance are covered in Core Skills training and are available from the website below:*

Links:  <https://turasdashboard.nes.nhs.scot>. Please complete the registration process (if you are not already registered) and search "core skills".

Managing complex cases

Some people will be experiencing such harm and impact from cocaine use that they are unable to undertake this work and may have poor engagement with addiction services. These people should be appropriately RAG and CRAFT risk assessed. They may require assertive outreach, crisis or stabilisation residential care where available, or an inpatient admission if they have mental health complexity. People experiencing harm from primary cocaine or poly drug use could be discussed in supervision and then possibly at MDT depending on the level of complexity and harm.

There are a range of residential services that support individuals at varying stages of their recovery. Resource and availability may be different in each health board area.

Bibliography

Adult Mental Health & Addictions Service Shared Guidance and Specification for Interface Working March 2021. <https://rightdecisions.scot.nhs.uk/media/cwjefaho/mhs-03-adult-mental-health-and-addictions-services-shared-guidance-and-specification-for-interface-working.pdf>

Crew.scot. 2022. *Crew 2000 – Mind altering*. [online] Available at: <https://www.crew.scot/> [Accessed 10 February 2022].

Darke, S., Kaye, S., McKetin, R. and Duflou, J., 2007. Physical and psychological harms of psychostimulant use. *National Drug and Alcohol Research Center*.

Drug Related Deaths in Scotland in 2020. National Records of Scotland. 30th July 2021. <https://www.nrscotland.gov.uk/files//statistics/drug-related-deaths/20/drug-related-deaths-20-pub.pdf>

Fitzpatrick, J., 2022. *Shared Protocol for Learning Disability & Mental Health Interface Working*.

Havakuk, O., Rezkalla, S.H. and Kloner, R.A., 2017. The cardiovascular effects of cocaine. *Journal of the American College of Cardiology*, 70(1), pp.101-113.

Network, N.P.T.U., 2015. Guidance on the clinical management of acute and chronic harms of club drugs and novel psychoactive substances. *NEPTUNE, London*.

NHS Lothian Quality Prescribing Group Substance Misuse Directorate Good Practice Guidance: Prescribing Opiate Substitution Therapy (OST) for Opiate Dependent Individuals also Using Stimulants A Fletcher, A Hart, M Kehoe, C Miller, I Scott, J Smith May 2019.

Perper, J.A. and Thiel, D.H.V., 1992. Respiratory complications of cocaine abuse. *Recent Developments in Alcoholism*, pp.363-377.

Scottish Advisory Committee on Drug Misuse. Psychostimulant Working Group. Substance Misuse Division, 2002. *Scottish Advisory Committee on Drug Misuse: Psychostimulant Working Group Report*. Scottish Executive.

Strang, J. and Independent Expert Working Group, 2017. Drug misuse and dependence: UK guidelines on clinical management.

Appendices

Cocaine Assessment Tool

Which drugs have you taken in the last 6 months and how often do you typically take them?					
	Less than once a week	Weekly	Most days	Daily	More than once a day (how many times?)
Heroin					
Methadone (street)					
Methadone (prescribed)					
Buprenorphine (espranor/ subutex) and/or Buprenorphine + Naloxone (suboxone)(street)					
Buprenorphine (espranor/ subutex)and/or Buprenorphine + Naloxone (suboxone) (prescribed)					
Benzodiazepine(street)					
Benzodiazepine (prescribed)					
Alcohol					
Pregabalin (prescribed)					
Pregabalin (street)					
Gabapentin (prescribed)					
Gabapentin (street)					
Cocaine powder					
Cocaine freebase/crack					
Cocaine and heroin (snowball)					
Amphetamine					
Methamphetamine					
New psychoactive substance stimulant type (NPS)					
Steroid (IPEDS)					
Human Growth Hormone (IPEDS)					
Tanning agents/melanotan (IPEDS)					
Other (what?):					
Are you on any other prescribed medications? (please specify): e.g. ARV, anti-depressants etc.					

Patterns of cocaine/crack use

How would you describe your use?	
Occasional use	
Regular use	
Binge use	
Daily use	
Over the last 6 months on average how much would you typically use in a day/ drug using session?	
If you binge, how many days would be a typical binge session?	
During the past 6 months, how often have you had a strong desire or urge to use cocaine/crack?	
Never/almost never	
Sometimes	
Often	
Always/almost always	
How long does a single hit last for?	
Less than 15 mins	
15-30mins	
30 mins-1 hour	
1-2 hours	
2-4 hours	
How long do you usually leave between hits?	
Less than 30 mins	
30-60mins	
1-2 hours	
2-4 hours	
4-6 hours	
6-8 hours	
Over the last 6 months on average how many hits do you get out of a gram/rock etc.?	

Dependence

Over the last 3 months: Did you ever think your use of cocaine was out of control?			
Never/almost never	Sometimes	Often	Always/nearly always
Did the prospect of missing a hit make you very anxious or worried?			
Never/almost never	Sometimes	Often	Always/nearly always
Did you worry about your use of cocaine?			
Never/almost never	Sometimes	Often	Always/nearly always
Did you wish you could stop using cocaine?			
Never/almost never	Sometimes	Often	Always/nearly always
How difficult would you find it to stop or go without?			
Not difficult	Quite difficult	Very difficult	Impossible

Side effects

How often do you experience any of the following when you use cocaine/crack?				
	Never/almost never	Sometimes	Often	Always/nearly always
Chest pain				
Palpitations/racing heart				
Fits/seizures				
Pain or burning at injecting site				
Coughing				
Difficulty breathing				
Burns/blisters/sores				
Disturbed sleep/insomnia				
Agitation				
Fear or panic				
Anxious				
Confused				
Paranoid				
Hear or see things				
Thinking people are trying to harm you or you are being watched/bugged etc.				
Thoughts of harming yourself or others				
Overdose				
Comedown effects, e.g. irritability, low mood				

Physical/Mental health history

Any current or previous health history which may impact on your cocaine/crack use?
E.g. physical health issues or concerns / attend hospital consultants etc.? history of contact with mental health services or treatment from GP

Poly substance use

Do you take other substances to help manage any of the side effects you experience?
(tick all that apply)

Heroin	
Benzodiazepines	
Sleeping tablets	
Alcohol	
Cannabis	
Other (what?):	

Setting

Where do you typically use? (tick all that apply)

Own home	
Friends home	
Shelter	
Hostel	
Safe consumption facility	
Prison	
Public spaces	
Other (where?):	

Do you use alone?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Hygiene

Do you clean your hands prior to preparing drugs?	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
How do you clean them?	
Warm water and soap	
Alcohol type hand cleanser	
Cold water and soap	
Warm water no soap	
Cold water no soap	
Other (please specify):	

Sexual Health and BBVs

When using cocaine/crack do you have unplanned or impulsive sex?	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
When using cocaine/crack have you had sex you later regretted?	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
When using cocaine/crack do you have rough sex? E.g. vigorous intercourse, fisting etc.	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
When using cocaine/crack do you have unprotected sex? (worker should explore where relevant all forms of sex including oral, vaginal and sharing of sex toys)	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
Are you pregnant? Is it possible you could be pregnant?	

Have you had any sexual relationships in order to get things you need e.g. money, food, alcohol or drugs, gifts or other things that are important to you?	
Have you been tested for HIV in last six months?	
Are you aware of current HIV status? i.e. collected results and no risk exposure since test	
Have you been tested for HCV in last six months?	
Are you aware of current HCV status? i.e. collected results and no risk exposure since test	
Have you had a sexual health check up in the last six months?	
Are you aware of current STI status? i.e. collected results and no risk exposure since test	

Injecting (please only complete this section if client injects)

How do you inject?				
	Cocaine	Heroin	Amphetamine	Other (what?):
IV				
IM				
Subcutaneous				

What size needle do you generally use? (please tick)

Colour	Guage Size	Length	Suitability for	Drugs usually injected by needle	Available in One Hit Kit Format
Green	21g (0.8mm)	1 1/2" (38mm)	Drawing oil based steroids. Intramuscular (IM) Injection of steroids (buttocks).	Oil Based Steroids	In Steroid Glasses Case
Blue	23g (0.6mm)	1 1/4" (32mm)	Intramuscular (IM) injection of steroids (buttocks). Femoral (groin) injection (IV).	Steroids, Heroin, Cocaine and Amphetamine (if femoral vein accessed).	In Blue One Hit Kit (2ml) In Steroid Glasses Case
Blue	23g (0.6mm)	1" (25mm)	Intramuscular (IM) injection (buttock, thighs and shoulders) of steroids. Femoral (groin) injection (IV).	Steroids, Heroin, Cocaine and Amphetamine (if femoral vein accessed).	No-Access fixed site for Pick & Mix
Orange	25g (0.5mm)	1" (25mm)	Femoral (groin) injection (IV). Slightly deeper veins when the needle is prone to blocking.	Steroids, Heroin, Cocaine and Amphetamine (if femoral vein accessed).	In Orange One Hit Kit (3ml)
Orange	25g (0.5mm)	5/8" (16mm)	Slightly deeper veins when the needle is prone to blocking.	Steroids, Heroin, Cocaine and Amphetamine (if femoral vein accessed).	No-Access fixed site for Pick & Mix
1ML Fixed (LOW DEAD SPACE)	27g-29g	1/2" (13mm)	Superficial veins, such as arms, hands, feet and legs (IV) Subcutaneous injection of some hormones.	Heroin, Cocaine, Amphetamine and NPS. IPEDS such as growth hormone, tanning agents and peptides.	In Black One Hit Kit (1ml)

Over the past 6 months, which of the following sites have you used for injecting?

	Cocaine	Heroin	Amphetamine	Other (what?):
Arms				
Hands				
Feet				
Leg				
Groin				
Neck				
Breasts				
Penis				
Other				

How frequently do you inject?

Less than once a week	
Weekly	
Most days	
Daily	
More than once a day (how many times)?	

Over the past 6 months have you experienced any complications which may be connected to you injecting?

Abscesses	
infection	
Cellulitis	
Ulcer	
DVT	
Open Wound	
Blocked or Collapsed Veins	
Missed hits	
Other (what?):	

Do you ever flush the blood back and forth whilst the needle is still in the vein?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Do you ever use any of the following techniques for raising a vein?

Tourniquet	
Warm water	
Gentle exercise	
Fist clenching	
Swinging arms(windmill motion)	

Preparation for injecting

Do you clean your injecting sites prior to injecting drugs?	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
How do you clean them?	
Warm water and soap	
Alcohol type hand cleanser	
Cold water and soap	
Warm water no soap	
Cold water no soap	
Other (please specify):	

Use of acidifier

Do you use an acidifier to break down your drugs for injection?	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
If yes how much do you tend to use?	
Whole content of single use sachet	
2 or more single use sachets	
Only a small amount of sachet	
Other (please specify):	

Use of water

What source of water do you use to prepare your drug for injection?	
Water from injection ampules	
Water from cooled boiled kettle	
Cold tap water	
Warm or hot water from tap	
Bottled water	
Water from a cup that others have access to	
Other (please specify):	

Do you use water to flush out your syringe after injecting?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Needle reuse and batch preparation

Do you ever share needles/syringes with anyone else?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Do you ever share spoons, water, filter with anyone else?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Do you ever reuse your own needle/syringe?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Do you ever reuse your own spoons, water, filter?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Do you ever prepare (for injecting) a larger amount of drugs to share with others?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Do you ever have your drugs for injection prepared by others?

Never/almost never	
Sometimes	
Often	
Always/nearly always	

Smoking (please only complete this section if client inhales)

What kind of inhalation pipes and gauzes do you use?	
Glass/Pyrex pipe	
Steel pipe	
Foil from IEP service	
Pipe made from drink can	
Pipe made from plastic bottle	
Pipe made from inhaler	
Pipe made from glass miniature bottle	
Pipe gauze	
Gauze made from metal scouring pad	
Gauze made from tin foil	
Other (please specify):	
Do you ever share inhalation pipes with others?	
Never/almost never	
Sometimes	
Often	
Always/nearly always	
How often do you change your gauze?	
Every hit/almost every hit	
Every few hits	
Occasionally	
Rarely	
Never	
When using crack/freebase do you? (tick all that apply)	
Break up the rock	
Hold the smoke in the lungs	
Scrape down residue from pipe	
Smoke with tobacco/ash	
Allow pipe to cool to touch	

Freebase cocaine

If you smoke freebase cocaine, do you typically:

Buy as freebase

Make your own freebase

If you make your own freebase do you rinse off the ammonia and dry thoroughly before smoking?

Never/almost never

Sometimes

Often

Always/nearly always

Motivational Enhancement: Identifying Problems

Below are some problems that can be made worse by drug and alcohol use.

Put a check beside any that you have had.

Medical or Physical:

- Head Injury in past
- Overdose experiences
- Stomach problems
- Dental/teeth problems
- Seizures or convulsions
- Large weight gain or loss
- Diarrhoea or constipation
- Nose or sinus problems
- HIV issues
- Hepatitis issues
- Waking up at night with a start
- Difficulty breathing
- Difficulty breathing at night
- Heart problems
- Chronic pain
- Specific pain problem
- Chronic fatigue
- Memory problems
- Other (Specify):

Relationships:

- Fights with partner
- Fights with children
- Fights with other family or friends
- Feeling alone
- Difficult to talk to other people
- Difficulty solving problems
- Loss of friends
- Only knowing people who use
- Problems with children/parenting
- Loss of partner
- Children at risk/in care
- Problems in sex life
- Other (Specify):

Legal:

- Arrested - possession/dealing
- Arrested - theft or robbery
- Arrested - assault/other violence
- DTTO, probation or parole
- Divorce or separation
- Child visitation issues
- Other (Specify):

Emotions and Feelings:

- Depression
- General anxiety or stress
- Panic attacks
- Anxiety around other people
- Anxiety when outside
- Other specific fears/phobias
- Sudden swings in mood
- Problems controlling anger
- Problems dealing with the past
- Remembering/flashbacks to past
- Hallucinations
- Feeling suspicious or paranoid
- Memory problems
- Can't sit still - always moving
- Can't relax
- Can't concentrate
- Other (Specify):

Housing, Finances, and Skills:

- Finding a place to live/sleep
- Furnishing/equipping your home
- Finances and budgeting
- Home skills (cooking, shopping)
- Taking care of your home
- Taking care of yourself
- Other (Specify):

Daily Routines

- Problems getting to sleep
- Severe snoring
- Problems waking up
- Eating too much or too little
- Not eating a balanced diet
- Bored during the day
- Lack of fun things to do
- Loss of sports or hobbies
- Lack of physical exercise
- Problems finding or keeping work
- Problems with training or school
- Other (Specify):

Laura Freeman, Ph.D. Addiction Interventions

Motivational Enhancement: Identifying Things to Work On

Now, look at this list again.

- 1) Check what would you like to change in your life.
- 2) Number those checked in order of importance.

Medical or Physical:

- Head Injury in past (#:)
- Overdose experiences
- Stomach problems
- Dental/teeth problems
- Seizures or convulsions
- Large weight gain or loss
- Diarrhoea or constipation
- Nose or sinus problems
- HIV issues
- Hepatitis issues
- Waking up at night with a start
- Difficulty breathing
- Difficulty breathing at night
- Heart problems
- Chronic pain
- Specific pain problem
- Chronic fatigue
- Memory problems
- Other (Specify):

Relationships:

- Fights with partner
- Fights with children
- Fights with other family or friends
- Feeling alone
- Difficult to talk to other people
- Difficulty solving problems
- Loss of friends
- Only knowing people who use
- Problems with children/parenting
- Loss of partner
- Children at risk/in care
- Problems in sex life
- Other (Specify):

Legal:

- Arrested – possession/dealing
- Arrested - theft or robbery
- Arrested - assault/other violence
- DTTO, probation or parole
- Divorce or separation
- Child visitation issues
- Other (Specify):

Emotions and Feelings:

- Depression
- General anxiety or stress
- Panic attacks
- Anxiety around other people
- Anxiety when outside
- Other specific fears/phobias
- Sudden swings in mood
- Problems controlling anger
- Problems dealing with the past
- Remembering/flashbacks to past
- Hallucinations
- Feeling suspicious or paranoid
- Memory problems
- Can't sit still – always moving
- Can't relax
- Can't concentrate
- Other (Specify):

Housing, Finances, and Skills:

- Finding a place to live/sleep
- Furnishing/equipping your home
- Finances and budgeting
- Home skills (cooking, shopping)
- Taking care of your home
- Taking care of yourself
- Other (Specify):

Daily Routines

- Problems getting to sleep
- Severe snoring
- Problems waking up
- Eating too much or too little
- Not eating a balanced diet
- Bored during the day
- Lack of fun things to do
- Loss of sports or hobbies
- Lack of physical exercise
- Problems finding or keeping work
- Problems with training or school
- Other (Specify):

Laura Freeman, Ph.D. Addiction Interventions

Motivational Enhancement: Daily Routine Worksheet

The purpose of this worksheet is to provide participants with the opportunity to relate the content of the course with their work. The following questions can be used as a personal review of the material, as a part of a consultation with other professionals, or a structure for peer or individual supervision. Consider an individual with whom you work, either currently or in the past....

1. How do they describe their mental health in general? How do they describe their sleep? What are their beliefs about their mental health, and what do they tell themselves?

2. What time do they wake up in the morning and what do they do? What are their morning rituals?

3. What types of physical exercise do they get during the day? Do they get outside in the sunshine? What about mental exercise?

4. What do they eat and when?

5. List all the substances they use, including legal, illicit and prescribed.

6. Which parts of their living space do they use? Do they spend most of their time in one place? Do they use their bedroom for “active” activities while awake?

7. What do they do in the 2 hours before bedtime? What are their “wind down” rituals?

8. What time do they go to sleep? What do they do when if they are unable to fall asleep?

9. How do they sleep during the night? Do they describe waking up with a start? Nightmares? What do they do to cope with any sleep problems?

Motivational Enhancement: Happiness Scale

This scale is intended to estimate your *current* happiness with your life in each of the ten areas listed below. Ask yourself the following question as you rate each area:

How happy am I with this area of my life?

You are to circle one of the numbers (1-10) beside each area.

Numbers toward the left indicate various degrees of unhappiness, while numbers toward the right reflect various levels of happiness.

In other words, state according to the numerical scale (1-10) exactly how you feel today.

Remember: Try to exclude all feelings of yesterday and concentrate only on the feelings of today in each of the life areas. Also try not to allow one category to influence the results of the other categories.

	Completely Unhappy					Completely Happy				
	1	2	3	4	5	6	7	8	9	10
Drug use	1	2	3	4	5	6	7	8	9	10
Job or Education Progress	1	2	3	4	5	6	7	8	9	10
Money Management	1	2	3	4	5	6	7	8	9	10
Social Life	1	2	3	4	5	6	7	8	9	10
Personal Habits	1	2	3	4	5	6	7	8	9	10
Marriage/Family Relationships	1	2	3	4	5	6	7	8	9	10
Legal Issues	1	2	3	4	5	6	7	8	9	10
Emotional Life	1	2	3	4	5	6	7	8	9	10
Communication	1	2	3	4	5	6	7	8	9	10
General Happiness	1	2	3	4	5	6	7	8	9	10

Adapted from: Meyers, R.J., and Smith, J.E. (1995) *Clinical Guide to Alcohol Treatment: The Community Reinforcement Approach*. New York: Guilford Press, p.95. Permission for use granted by Guilford Press. For the most recent overview of CRA, see: Meyers, R. J., Roozen, H.G., and Smith, J. E. (2011). The Community Reinforcement Approach: An Update of the Evidence. *Alcohol Research & Health*, 33 (4), 380-387.

Motivational Enhancement: Exploring the Relationships between Substance Use and Mental Health

This worksheet is designed to explore the relationship between substance use and mental health. First, list all of the legal, illicit, and prescribed substances currently being used, either regularly or periodically. Second, list the current mental health and social problems. Then, look for ways in which the two columns interact.

Substances Currently Being Used	Mental Health and Social Problems

What is the perceived coping potential of the substances used (e.g substance related beliefs?)

What is the impact of the substance use on mental health?

Motivational Enhancement: What I Want From Treatment:

Instructions

People have different ideas about what they want, need, and expect from treatment.

This questionnaire is designed to help you explain what you would *like* to have happen in your treatment. Many possibilities are listed. For each one, please indicate how much you would like for this to be part of your treatment.

You can do this by circling one number (0, 1, 2, or 3) for each item. This is what the numbers mean:

0 NO means that you definitely do **NOT** want or need this from treatment.

1 ? means that you are **UNSURE**. **MAYBE** you want this from treatment.

2 YES means that you **DO** want or need this from treatment.

3 YES! means that you **DEFINITELY** want or need this from treatment.

FOR EXAMPLE:

Consider item #1, which says, "I want to receive detoxification." If you definitely do NOT want or need to receive detoxification, you would circle **0**. If you are UNSURE whether you want or need detoxification, you would circle **1**. If you DO want detoxification, you would circle **2**. If you DEFINITELY know that detoxification is an important goal for your treatment, you would circle **3**.

If you have any questions about how to use this questionnaire, ask for assistance before you begin.

What I Want From Treatment

 DO YOU WANT THIS FROM TREATMENT?		No 0	Maybe 1	Yes 2	Yes!! 3
1.	I want to receive detoxification, to ease my withdrawal from alcohol or other drugs.	0	1	2	3
2.	I want to find out for sure whether I have a problem with alcohol or other drugs.	0	1	2	3
3.	I want help to stop drinking alcohol completely.	0	1	2	3
4.	I want help to decrease my drinking.	0	1	2	3
5.	I want help to stop using drugs (other than alcohol).	0	1	2	3
6.	I want to stop using tobacco.	0	1	2	3
7.	I want to decrease my use of tobacco.	0	1	2	3
8.	I want help with an eating problem.	0	1	2	3
9.	I want help with a gambling problem.	0	1	2	3
10.	I want to take Antabuse (a medication to help me stop drinking).	0	1	2	3
11.	I want to take a medication to help me stop using alcohol or heroin.	0	1	2	3

12.	I want to take methadone.	0	1	2	3
13.	I want to learn more about alcohol/drug problems.	0	1	2	3
14.	I want to learn some skills to keep from returning to alcohol or other drugs.	0	1	2	3
15.	I would like to learn more about self-help groups: 12-Step programs like Alcoholics Anonymous (AA) or Narcotics Anonymous (NA) or Smart Recovery.	0	1	2	3
16.	I would like to talk about some personal problems.	0	1	2	3
17.	I need to fulfil a requirement of the courts.	0	1	2	3
18.	I would like help with problems in my marriage or close relationship.	0	1	2	3
19.	I want help with some health problems.	0	1	2	3
20.	I want help to decrease my stress and tension.	0	1	2	3
21.	I would like to improve my health by learning more about nutrition and exercise.	0	1	2	3
22.	I want help with depression or moodiness.	0	1	2	3
23.	I want to work on my spiritual growth.	0	1	2	3
24.	I want to learn how to solve problems in my life.	0	1	2	3
25.	I want help with angry feelings and how I express them.	0	1	2	3
26.	I want to have healthier relationships.	0	1	2	3
27.	I would like to discuss sexual problems.	0	1	2	3
28.	I want to learn how to express my feelings in a more healthy way.	0	1	2	3
29.	I want to learn how to relax better.	0	1	2	3
30.	I want help in overcoming boredom.	0	1	2	3
31.	I want help with feelings of loneliness.	0	1	2	3
32.	I want to discuss having been physically abused.	0	1	2	3
33.	I want help to prevent violence at home.	0	1	2	3
34.	I want to discuss having been sexually abused.	0	1	2	3
35.	I want to work on having better self-esteem.	0	1	2	3
36.	I want help with sleep problems.	0	1	2	3

37.	I want help with legal problems.	0	1	2	3
38.	I want advice about financial problems.	0	1	2	3
39.	I would like help in finding a place to live.	0	1	2	3
40.	I could use help in finding a job.	0	1	2	3
41.	Someone close to me has died or left, and I would like to talk about it.	0	1	2	3
42.	I have thoughts about suicide, and I would like to discuss this.	0	1	2	3
43.	I want help with personal fears and anxieties.	0	1	2	3
44.	I want help to be a better parent.	0	1	2	3
45.	I feel very confused and would like help with this.	0	1	2	3
46.	I would like information about or testing for HIV/AIDS or Hepatitis C.	0	1	2	3
47.	I want someone to listen to me.	0	1	2	3
48.	I want to learn to have fun without drugs or alcohol.	0	1	2	3
49.	I want someone to tell me what to do.	0	1	2	3
50.	I want help in setting goals and priorities in my life.	0	1	2	3
51.	I would like to learn how to manage my time better.	0	1	2	3
52.	I want help to receive disability payments.	0	1	2	3
53.	I want to find enjoyable ways to spend my free time.	0	1	2	3
54.	I want help in getting my child(ren) back.	0	1	2	3
55.	I would like to talk about my past.	0	1	2	3
56.	I need help in getting motivated to change.	0	1	2	3
57.	I would like to see a female counsellor.	0	1	2	3
58.	I would like to see a male counsellor.	0	1	2	3
59.	I would like to see the counsellor I had before.	0	1	2	3
60.	I would like to see a doctor or nurse about medical problems.	0	1	2	3
61.	I want to receive medication.	0	1	2	3

62.	I would like my spouse or partner to be in treatment with me.	0	1	2	3
63.	I would like to have private, individual counselling.	0	1	2	3
64.	I would like to be in a group with people who are dealing with problems similar to my own.	0	1	2	3
65.	I need someone to care for my children while I am in treatment.	0	1	2	3
66.	I want my treatment to be short.	0	1	2	3
67.	I believe I will need to be in treatment for a long time.	0	1	2	3

Adapted from: William R. Miller & Janice M. Brown. Center for Substance Abuse Treatment. *Enhancing Motivation for Change in Substance Abuse Treatment*. Treatment Improvement Protocol (TIP) Series 35. DHHS Publication No. (SMA) 99-3354. Rockville, MD: Substance Abuse and Mental Health Services Administration, 1999.

Motivational Enhancement: Exploring the Pros and Cons of Making a Change

Beside each point, write down a number between 1 (Not Important) to 10 (Extremely) for how much it matters to you.

Making A Change: Positives	Making A Change: Negatives
Staying the Same: Positives	Staying the Same: Negatives

Laura Freeman, Ph.D. Addiction Interventions

Developing Control: Functional Analysis

This collaborative exercise helps to identify potential skills for managing relapse and developing an alternative lifestyle. Consider the identified problem behaviour. This may be substance use (e.g. alcohol binge) or mental health (eg. self-harm). Then consider what happens before the behaviour happens. Finally, explore the short and long term consequences.

Environmental Triggers	Thoughts and Feelings	Behaviour	Positive Consequences	Negative Consequences
			Short term: Long term:	Short term: Long term:

This exercise can help a person explore the relationship between triggers, their behaviour and the consequences. It is also useful in identifying skills needed to help the person manage triggers and control impulsive behaviour. This form can also be used to explore and reinforce alternative behaviours

Laura Freeman, Ph.D. Addiction Interventions

Developing Control: Functional Analysis of Recovery-Orientated Behaviour

This collaborative exercise explores healthy, recovery-orientated behaviours in your life. Choose something healthy or fun that you do right now that does not involve alcohol and/or drugs. Then consider what happens before the behaviour happens. Finally, explore the short and long term consequences.

Environmental Triggers (Who with? Where? When?)	Thoughts and Feelings (What were you thinking? Feeling emotionally? Feeling physically?)	Behaviour (Be specific – What do you do, how long and how often?)	Positive Consequences (Relationship? Emotions? Physical Health? Legal? Job/education? Money?)	Negative Consequences (What are the barriers? Who, where, when? Thoughts, emotions, physical?)
			Short and long term:	Short and long term:

Laura Freeman, Ph.D. Addiction Interventions

Developing Control: Discovering Triggers

Discovering Triggers of Your Alcohol and Other drugs Use

1. List the places where you are **most likely** to use alcohol and other drugs:

2. List the people with whom you are **most likely** to use alcohol and other drugs:

3. List and times or days when you are **more likely** to use alcohol and other drugs:

4. List any activities that make it more likely that you will use alcohol and other drugs:

5. Do you think that you use alcohol and other drugs when you are feeling certain ways? Read through the following list and mark the ones that are relevant to you. For those you have marked, list specific examples from your own experience.

- a. At the end of (or during) a tense day
- b. When faced with something you fear or are anxious about
- c. When you've failed to accomplish something you'd planned
- d. When you feel you have been taken advantage of
- e. When you are bored
- f. When you are in a social situation
- g. When you feel bad about yourself
- h. When you are depressed
- i. When you want to feel energized or high
- j. When you are faced with a tough problem
- K. When you want to be friendly
- l. When you wish your personality was different
- m.others not listed here

6. List the places where you are **unlikely** to use alcohol and other drugs:

7. List the people with whom you are **unlikely** to use alcohol and other drugs:

8. List the times or days when you are **unlikely** to use alcohol and other drugs:

9. List the activities you engage in when you are **unlikely** to use alcohol and other drugs:

Adapted from: Budney, A. and Higgins, S. (1998). *A Community Reinforcement Plus Vouchers Approach: Treating Alcohol and other drugs Addiction*. NIDA. Page 60. This form was originally adapted from Miller and Munoz 1982

Developing Control: Self-Management Plan

Self-Management Planning Sheet

Trigger	Plans	+/- Consequences	Difficulty (1 – 10)
1.			

Adapted from: Budney, A. and Higgins, S. (1998). *A Community Reinforcement Plus Vouchers Approach: Treating Alcohol and other drugs Addiction*. NIDA. Page 60. This form was originally adapted from Miller and Munoz 1982

Developing Control: Coping with Cravings and Urges

Coping With Cravings and Urges

Reminders:

- Urges are common and normal. They are not a sign of failure. Instead, try to learn from them about what your craving triggers are.
- Urges are like ocean waves. They get stronger only to a point, then they start to go away.
- If you don't use, your urges will weaken and eventually go away. Urges only get stronger if you give in to them.
- You can try to avoid urges by avoiding or eliminating the cues that trigger them.
- You can *cope* with urges by -
 - » Distracting yourself for a few minutes.
 - » Talking about the urge with someone supportive.
 - » "Urge surfing" or riding out the urge.
 - » Recalling the negative consequences of using. - Talking yourself through the urge.

Each day this week, fill out a daily record of craving and what you did to cope with craving.

Example:

Date/Time	Situation, thoughts, and feelings	Intensity of Craving (1-100)	Length of Craving	How I Coped
Friday, 3 pm	Fight with boss, frustrated, angry	75	20 minutes	Called home, talked to Mary
Friday, 7 pm	Watching TV, bored, trouble staying awake	60	25 minutes	Rode it out and went to bed early
Saturday, 9 pm	Wanted to go out and get a drink	80	45 minutes	Played basketball instead

Developing Control: Coping with Cravings and Urges

Date/Time	Situation, thoughts, and feelings	Intensity of Craving (1-100)	Length of Craving	How I Coped

Adapted from: Carroll, K. (1998). A Cognitive-Behavioral Approach: Treating Cocaine Addiction. NIDA. Pages 53-54. This form was originally adapted from Kadden et al. 1992.

Developing Control: Coping with Thoughts about Using

Coping With Thoughts About Using

There are several ways of coping with thoughts about using alcohol and other drugs:

- Thinking through and remembering the end of the last time you used
- Challenging your thoughts
- Recalling the negative consequences of using
- Distracting yourself
- Talking through the thought

Before the next session, keep track of your automatic thoughts about using when they occur, and then record a positive thought and coping skills.

Thought about using substances	The positive thought or the coping skills used

Adapted from: Carroll, K. (1998). A Cognitive-Behavioral Approach: Treating Cocaine Addiction. NIDA. Page 64. This form was originally adapted from Monti et al. 1989.

Developing Control: Problem-solving Worksheet

Problem-solving Worksheet

Procedure:

1. *Gather information:* Recognize that a problem exists. Is there a problem? You get clues from your body, thoughts, feelings, behaviour, reactions to other people, and the ways that other people react to you. Think about the problem situation. Who is involved? When does it happen? Exactly what takes place? What effect does this have on you?
2. *Define the problem:* Describe the problem as accurately as you can. What goal would you like to achieve? Be as specific as possible. Break it down into manageable parts.
3. *Brainstorm for alternatives:* List all the things that a person in your situation could possibly do. Consider various approaches to solving the problem. Even list alternatives that seem impractical. Try taking a different point of view, try to think of solutions that worked before, and ask other people what worked for them in similar situations.
4. *Consider the consequences:* Look at each of your alternatives in turn. What things could you reasonably expect to result from taking each action? What positive consequences? What negative consequences are long-term? Which are short-term? Which do you think you could actually do?
5. *Make a decision:* Which alternative is the most likely to achieve your goal? Select the one likely to solve the problem with the least hassle. *Do it!* The best plan in the world is useless if it isn't put into action. Try it out.
6. *Evaluate its effectiveness:* Which parts worked best? Reward yourself for them. Would you do anything differently next time? After you have given the approach a fair trial, does it seem to be working out? If not, consider what you can do to strengthen the plan or give it up and try one of the other possible approaches. Remember that when you've done your best, you have done all you can do.

Continued on next page

Developing Control: Problem-solving Worksheet, Page 2

Practice Exercise

Choose a problem that may arise in the near future. Describe it as accurately as you can. Brainstorm possible solutions. Evaluate the potential outcomes. Prioritise solutions.

Identify the problem situation:

Brainstorm a list of possible solutions: Pros: Cons:

Developing Control: Seemingly Irrelevant Decisions

Seemingly Irrelevant Decisions

When making any decision, whether large or small, do the following:

- Consider all the options you have.
- Think about all the consequences, both positive and negative, for each of the options.
- Select one of the options. Pick a safe decision that minimizes your risk of relapse.
- Watch for “red flag” thinking - thoughts like “I have to . . .”, or “I can handle . . .” or
- “It really doesn’t matter if . . .”

Practice monitoring decisions that you face in the course of a day, both large and small, and consider safe and risky alternatives for each.

Decision	Safe alternative	Risky alternative

Adapted from: Carroll, K. (1998). A Cognitive-Behavioral Approach: Treating Cocaine Addiction. NIDA. Pages 76. This form was originally adapted from Monti et al. 1989.

Developing a Rewarding Lifestyle: Recreational Survey

Name: _____ Date: _____

Please tick the column which best describes your thoughts/feelings about each row.

Sports and Games				
Activity	Did at least 4x's last year	Would like to try...	Have tried and liked it...	No Interest
Football				
Tennis				
Swimming				
Golf				
Jogging/Running				
Aerobics				
Gymnastics				
Bowling				
Ten-pin bowling				
Darts				
Karate				
Pilates				
Tai Chi				
Yoga				
Snooker				
Billiards/Pool				
Rugby				
Badminton				
Sudoku				
Crossword Puzzles				
Internet Based Games				
Computer Games				
Chess				
Dominoes				
Scrabble				
Jigsaw Puzzles				
Table Tennis				
Shinty:				
Other:				

Outdoor Activities				
Activity	Did at least 4x's last year	Would like to try...	Have tried and liked it...	No Interest
Walking				
Hillwalking				
Bike/Dirt bike riding				
Fishing				
Going to the beach or shore				
Going to the park				
Ice skating				
Sailing				
Windsurfing				
Skateboarding				
Camping				
Gardening				
Picnics				
Birdwatching				
Other:				
Art, Music, Dance and Drama				
Drawing/Painting				
Photography				
Sculpture				
Woodworking				
Jewellery Making				
Knitting				
Sewing				
Cooking/Baking				
Listening to music				
Singing/Karaoke				
Playing instrument				
Dancing				
Scottish Country Dancing				
Line Dancing				
Acting				
Other:				

Other Activities				
Activity	Did at least 4x's last year	Would like to try...	Have tried and liked it...	No Interest
Shopping				
Museums				
Art Galleries				
Fairs				
Historical Sites and Events				
Sporting Events				
Eating Out				
Travelling				
Library				
Cinema				
Music Concers				
Theatre				
Talking with a Friend				
Attending a Party				
Volunteer Activities				
Watching TV				
Reading				
Meditation				
Creative Writing				
Support Groups				
Other:				
Other:				
Other:				

How would you rate your satisfaction with the recreational activities in your life?

Very Dissatisfied Dissatisfied Satisfied Very Satisfied

Are there problems that make it difficult for you to develop a healthy recreational lifestyle?
If yes, describe:

Developing a Rewarding Lifestyle: Goal Setting Worksheet for Reinforcing Recovery (With Directions)

Consider the Summary Worksheet. Select one or two of the areas, and decide on a goal(s) for the next month or so. (1) Find a goal that is positive. Say what you want to do - not what you don't want to happen. Make the goal brief, specific and decide on how you want to measure it. Make sure the goal is realistic and builds on things that you can do already. Next, what steps will you take to reach that goal (2) and how long will the step take (3)? Finally, what will you have as a reward after each step (4). This might be something good that comes out of the step, or you might plan a specific way to reward yourself positively and celebrate.

(1) Set a goal that is positive, brief, specific, measurable and realistic.	(2) What are the steps you plan to work towards your goal?	(3) How long will you plan for each step to take?	(4) <i>What will be the reward or "positive" with each step?</i>
Circle the general area you would like to make changes in: Alcohol and Other Drug Use Mental Health Physical Health Family + Significant Relationships Employment/Training/Education Recreational/Social Legal Goal for Change – I would like:	Step 1: Step 2: Step 3: Step 4: Step 5:	Step 1: Step 2: Step 3: Step 4: Step 5:	Step 1: Step 2: Step 3: Step 4: Step 5:

Developing a Rewarding Lifestyle: Brief Goal Setting Worksheet for Reinforcing Recovery

<p>(1) Set a goal that is positive, brief, specific, measurable and realistic.</p>	<p>(2) What are the steps you plan to work towards your goal?</p>	<p>(3) How long will you plan for each step to take?</p>	<p>(4) What will be the reward or "positive" with each step?</p>
<p>Circle the general area you would like to make changes in:</p> <p>Alcohol and Other Drug Use Mental Health Physical Health</p> <p>Family + Significant Relationships Employment/Training/Education</p> <p>Recreational/Social Legal</p> <p>Goal for Change – I would like:</p>	<p>Step 1:</p> <p>Step 2:</p> <p>Step 3:</p> <p>Step 4:</p> <p>Step 5:</p>	<p>Step 1:</p> <p>Step 2:</p> <p>Step 3:</p> <p>Step 4:</p> <p>Step 5:</p>	<p>Step 1:</p> <p>Step 2:</p> <p>Step 3:</p> <p>Step 4:</p> <p>Step 5:</p>


Laura Freeman, Ph.D. Addiction Interventions

Developing a Rewarding Lifestyle: Highlighting Progress Graph

Highlighting the progress made as someone works towards a goal is essential.

When someone is making changes in their lives, finding a visual way to see is often used to help people see the change happening. Highlighting progress can be as simple as a colour coded calendar, or as complex as graphs and charts. It is useful to create templates of charts, graphs or calendars which can be easily picked up and put to use.

Here are two different examples.



Laura Freeman, Ph.D. Addiction Interventions

Developing a Rewarding Lifestyle: Highlighting Progress Calendar

Name:		Month:				
<p>What do you want to see yourself doing? What's your goal? Choose a colour and use it to mark the days when you reach your current goal.</p>						
Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday

Laura Freeman, Ph.D. Addiction Interventions

Developing a Rewarding Lifestyle: Timeline Exercise

Major life events:

Birth Current

Timeline of mental health events:

Birth Current

Timeline of substance use:

Birth Current

Development of Beliefs and Assumptions

When (or at what periods) did problematic substance use happen?

Laura Freeman, Ph.D. Addiction Interventions

Where noted these worksheets have been supplied by the NHS Education for Scotland (NES) Core Behavioural and CBT Skills for Relapse Prevention and Recovery Management Course (Laura Freeman, NES, 2011). In order to follow and implement psychosocial interventions effectively, it is recommended that workers should have attended Core Skills training on TURAS.