

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	1 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

### Guidance Objective

To provide Healthcare Workers (HCWs) with details of the precautions necessary to minimise the risk of cross infection with Norovirus.

This guidance applies to all staff employed by NHS Greater Glasgow & Clyde and locum staff on fixed term contracts and volunteer staff.

This guidance applies to both confirmed and suspected outbreaks of Norovirus.

### KEY CHANGES FROM THE PREVIOUS VERSION OF THIS GUIDANCE

**Important Note:** The version of this policy found on the Infection Prevention & Control (eIPC Manual) on the intranet page is the only version that is controlled. Any other versions either printed or embedded into other documents or web pages should be viewed as uncontrolled and as such may not necessarily contain the latest updates, amendments, or linkages to other documents.

### Document Control Summary

Approved by and date	Board Infection Control Committee 15 <sup>th</sup> December 2022
Date of Publication	27 <sup>th</sup> January 2023
Developed by	Infection Prevention and Control Policy Sub-Group
Related Documents	National IPC Manual NHSGGC Hand Hygiene Guidance NHSGGC Loose Stools Guidance NHSGGC SOP Terminal Clean of Ward/Isolation Room NHSGGC SOP Twice Daily Clean of Isolation Rooms
Distribution/Availability	NHSGGC Infection Prevention and Control web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>
Lead Manager	Director Infection Prevention and Control
Responsible Director	Executive Director of Nursing

The most up-to-date version of this guidance can be viewed at the following web page:  
[www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control](http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control)


	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	2 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

## CONTENTS

Norovirus Aide Memoire .....	3
1. Responsibilities .....	4
2. General Information on Norovirus .....	5
3. Transmission Based Precautions for Norovirus.....	6
4. Evidence Base .....	9
Appendix 1 –Bowel Movement Record .....	10
Appendix 2 - Norovirus Outbreak Daily Checklist.....	11
Appendix 3 - Norovirus Outbreak Data Record Ward .....	12

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	3 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

### Norovirus Aide Memoire



	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	4 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

## 1. Responsibilities

### Healthcare Workers (HCWs) must:

- Follow this guidance.
- Inform their line manager if this guidance cannot be followed.
- Be vigilant for increased number of cases with similar symptoms and notify IPCT
- Implement care checklist

### Senior Charge Nurse (SCN) / Managers must:

- Support HCWs and Infection Prevention and Control Teams (IPCTs) in following this guidance.

### Infection Prevention and Control Teams (IPCTs) must:

- Keep this guidance up-to-date.
- Provide education opportunities on this guidance.
- Provide advice during outbreaks and incidents

### Occupational Health Service (OHS) must:

- Advise HCWs regarding possible infection exposure and return to work issues as necessary

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	5 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

## 2. General Information on Norovirus

<b>Communicable Disease/ Alert Organism</b>	<p>Norovirus is highly infectious and causes outbreaks of gastroenteritis in places where people congregate, e.g. schools, hospitals, nursing homes, cruise ships etc. Gastroenteritis caused by Norovirus is usually self-limiting, mild to moderate in severity and normally occurs during winter and early spring but can occur throughout the year. The infective dose is very small, between 10 – 100 virus particles. (HPS 2016).</p> <p>Norovirus outbreaks usually require closure of wards to prevent onward spread</p>
<b>Clinical Condition</b>	<p><b>Gastroenteritis:</b> Gastro-intestinal symptoms, e.g. nausea, vomiting (often projectile), non-bloody watery diarrhoea; characteristically lasting 12-48 hours. Also present may be abdominal cramps, myalgia (muscle pain), headache, malaise and low grade fever.</p>
<b>Mode of Spread</b>	<p><b>Direct Contact:</b></p> <ul style="list-style-type: none"> <li>• Hands come into contact with faecal matter/ vomit and subsequently touch the mouth.</li> </ul> <p><b>Indirect Contact:</b></p> <ul style="list-style-type: none"> <li>• Hands come into contact with contaminated equipment or contaminated surfaces and subsequently touch the mouth.</li> <li>• Consumption of faecal contamination in food or water.</li> </ul> <p><b>Droplet Dissemination:</b></p> <ul style="list-style-type: none"> <li>• Patients with excessive or projectile vomiting can disseminate large quantities of virus in droplets which can contaminate extensive areas of the ward/department.</li> </ul> <p><b>NB:</b> Norovirus can survive on any surface including equipment and on refrigerated food for up to 10 days.</p>
<b>Incubation period</b>	Usually 12-48 hours.
<b>Notifiable disease</b>	No.
<b>Period of communicability</b>	During the acute stage of the disease and up to 48 hours after symptoms of diarrhoea and vomiting have resolved or stools have returned to their normal (pre-infection pattern).
<b>Persons most at risk</b>	All. Susceptibility is widespread. It should be noted that mortality associated with Norovirus can occur particularly in elderly patients with co-morbidities. (HPS 2016)

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	6 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

### 3. Transmission Based Precautions for Norovirus

<b>Accommodation (Patient Placement)</b>	All patients symptomatic of unexplained vomiting and/or diarrhoea must be placed in a single room with en suite (or own dedicated commode). If the patient is clinically unsuitable for isolation, a risk assessment must be undertaken by the clinical team. Failure to isolate must be documented in the IPC Care Checklist and reviewed daily. If a single room is not available, after consulting bed manager, staff must inform a member of the IPCT to allow further assessment.
<b>IPC Care Checklist available</b>	Yes. Loose Stools Care Checklist
<b>Clinical / Healthcare Waste</b>	Waste should be designated as clinical/ healthcare waste and placed in an orange waste bag. Please refer to the <a href="#">NHSGCC Waste Management Policy</a>
<b>Domestic Advice</b>	<ul style="list-style-type: none"> <li>Patients room/bed space should be cleaned twice daily as per NHSGGC SOP <a href="#">Twice Daily Clean of Isolation Rooms</a></li> <li>Chlorine based detergents should be used for routine and terminal cleaning of the area.</li> <li>Blood and/ or body fluid contamination of the environment should be managed as per the NHSGGC Decontamination Guidance.</li> <li>On resolution of symptoms ( more than 48 hours asymptomatic) or discharge home, patient room/ bed space must be terminally cleaned see NHSGGC SOP <a href="#">Terminal Clean of Ward/Isolation Room</a></li> </ul>
<b>Equipment</b>	<p>Patient equipment must be dedicated as far possible, while patient remains symptomatic and during infectious period.</p> <p>Patient care equipment should be cleaned after use or twice daily with chlorine based detergent or immediately if visibly contaminated.</p> <p>Where possible equipment such as commodes, call buttons, washbowls, chairs, hoist slings, BP cuffs, thermometers etc. should be kept for use by individual, symptomatic patients. If equipment is taken out of the room it must be cleaned with 1000ppm chlorine based detergent.</p> <p>Please also refer to NHSGGC <a href="#">Decontamination Guidance</a></p>
<b>Hand Hygiene (HH)</b>	Hand hygiene is the single most important measure to prevent cross-infection with Norovirus.

The most up-to-date version of this guidance can be viewed at the following web page:  
[www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control](http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control)

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	7 of 12
	<p style="text-align: center;"><b>NOROVIRUS GUIDANCE TRANSMISSION BASED PRECAUTIONS</b></p>	Effective From	Dec 2022
		Review Date	Dec 2024
		Version	7
<p>The most up-to-date version of this guidance can be viewed at the following web page:  <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a></p>			
	<p>Hands must be decontaminated with liquid soap and water before and after each direct patient contact, after contact with the environment, after exposure to body fluids and before any aseptic tasks. Patients should be encouraged/ assisted to use the hand hygiene facilities after using the toilet or commode and before meals. Hand wipes should be provided to those patients unable to use hand hygiene facilities</p> <p>Visitors must also be encouraged to wash their hands with soap and water after visiting a patient with loose stools.</p> <p>Please refer to NHSGGC <a href="#">Hand Hygiene Guidance</a></p>		
<b>Individual precautions required until</b>	Patient has been asymptomatic for 48 hours.		
<b>Last Offices</b>	No special requirements.		
<b>Linen</b>	<p>Treat used linen as soiled/ infected, i.e. place in a water soluble bag, then into a clear plastic bag (brown bag used in Mental Health areas), tied, then into a white laundry hamper bag. Some wards and departments may use red laundry hamper bags as the outer bag.</p> <p>Please refer to <a href="#">National Guidance on the safe management of linen</a></p> <p>Any soiled clothing for home laundering should be placed into a domestic water soluble bag then into a patient clothing bag before being sent home. All soiled clothing for home laundering should be accompanied with a <a href="#">Washing Clothes at Home Leaflet</a> and staff should alert relatives / carers to the condition of the laundry. NB it should be recorded in the nursing notes that both advice and the information leaflet has been issued.</p>		
<b>Moving Patients between wards, hospitals and departments (including theatres)</b>	Movement of patients should be restricted until they have been asymptomatic for 48 hours. Movement of patients while symptomatic must only occur if there is a <b>clinical</b> need and this should be discussed with the IPCT and the receiving area.		
<b>Notice for Door</b>	The yellow IPC isolation sign must be placed on the door to the patient's room. In Mental Health Services, on advice of the IPCT. The door should remain closed and if the door cannot be closed,		

The most up-to-date version of this guidance can be viewed at the following web page:  
[www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control](http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control)

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	8 of 12
	<p style="text-align: center;"><b>NOROVIRUS GUIDANCE TRANSMISSION BASED PRECAUTIONS</b></p>	Effective From	Dec 2022
		Review Date	Dec 2024
		Version	7
<p>The most up-to-date version of this guidance can be viewed at the following web page:  <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a></p>			
	then an IPCT risk assessment should be completed and reviewed frequently.		
<b>Outbreak</b>	<p>If an outbreak is suspected, contact a member of the IPCT and refer to the NHSGGC <a href="#">Outbreak Incident Management Plan</a> for further guidance.  <b>Appendices 2 and 3 can be used to gather information during the outbreak.</b></p>		
<b>Personal Protective Equipment (PPE)</b>	Disposable yellow aprons and disposable gloves must be worn if in contact with an infected patient or their environment. Where there is a risk of blood and / or body fluid splash to the face, a fluid resistant surgical mask and eye protection must be considered.		
<b>Specimens required</b>	Specimens of faeces must be obtained for microbiology and virology at the earliest possible opportunity. Both faecal and vomit specimens can be sent to virology.		
<b>Stool Charts</b>	It is the responsibility of staff within the area to record type/frequency of stool, using the appropriate stool chart. See <a href="#">Appendix 1</a> Bowel Movement Record		
<b>Terminal Cleaning of Room</b>	Refer to <a href="#">Terminal Clean of Ward/Isolation Room SOP</a>		
<b>Visitors</b>	Visitors are not required to wear aprons and gloves, unless they are participating in patient care. Visitors must be advised to decontaminate their hands with liquid soap and water on leaving the room/ patient. Symptomatic visitors should be advised not to visit patients in hospital until they have been asymptomatic for 48 hours. Staff should consider restricting the number of visitors to two and advising visitors not to bring young children and babies to visit whilst the patient is symptomatic.		

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	9 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

#### 4. Evidence Base

HPS (2016) General information to prepare for and manage norovirus in care settings.  
[HPS Website - General information to prepare for and manage norovirus in care settings \(scot.nhs.uk\)](http://www.scot.nhs.uk/hps/prepare-for-and-manage-norovirus-in-care-settings)

CDC (2017) Guideline for the Prevention and Control of Norovirus Gastroenteritis Outbreaks in Healthcare Settings

CDC (2011) Updated Norovirus Outbreak Management and Disease Prevention Guidelines. MMWR 60(RR03); 1-15.

Chadwick P.R. et al (2000) Management of hospital outbreaks of gastro-enteritis due to small round structured viruses. Report of the Public Health Laboratory Service Viral GastroEnteritis Working Group. Journal of Hospital Infection. 45 pp 1-10

Heymann D.L. (2022) Control of Communicable Diseases Manual. 21<sup>st</sup> ed. The official report of the American Public Health Association. Washington DC.

[ARHAI \(2022\) National Infection prevention and Control Manual](http://www.cdc.gov/nipw/ARHAI-2022-National-Infection-prevention-and-Control-Manual)

The most up-to-date version of this guidance can be viewed at the following web page:  
[www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control](http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control)

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE	Page	10 of 12
		Effective From	Dec 2022
	<b>NOROVIRUS GUIDANCE</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Dec 2024
		Version	7
The most up-to-date version of this guidance can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

**Appendix 1 – Bowel Movement Record**

**BOWEL MOVEMENT RECORD**

Month: \_\_\_\_\_ Year: \_\_\_\_\_

Name: \_\_\_\_\_

Date	Time	Size S-small M-medium L-large S M L	Type 1	Type 2	Type 3	Type 4	Type 5	Type 6	Type 7	Staff Initials
			Separate hard lumps like nuts (hard to pass) 	Sausage shaped but lumpy 	Like a sausage but with cracks on surface 	Like a sausage or snake, smooth and soft 	Soft blobs with clear-cut edges (passed easily) 	Huffy pieces with ragged edges, a mushy stool 	Watery, no solid pieces (entirely liquid) 	
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									
	am									
	pm									

Adapted from the Bristol Stool Scale developed by KW Heaton and SJ Lewis at the University of Bristol, 1997

The most up-to-date version of this guidance can be viewed at the following web page:  
[www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control](http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control)

	NHS GREATER GLASGOW & CLYDE CONTROL OF INFECTION COMMITTEE POLICY	Page	11 of 12
		Effective From	Sept 2020
	<b>NOROVIRUS POLICY</b> <b>TRANSMISSION BASED PRECAUTIONS</b>	Review Date	Sept 2022
		Version	6
The most up-to-date version of this policy can be viewed at the following web page: <a href="http://www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control">www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control</a>			

### Appendix 2 - Norovirus Outbreak Daily Checklist

Both the checklist and data record to be completed and updated by the ward staff.

Norovirus Outbreak Daily Checklist to ensure Norovirus Control Measures are in place.

Date commenced: \_\_\_\_\_

Tick if done, X if not done, N/A for not applicable.

Hospital:	Ward:	ICT informed date:	Date:								
<b>The ward is closed</b> due to admissions and transfers – until <b>48</b> hours after last new case.											
<b>The ward (and side-room) doors are closed</b> and there is an approved notice on the ward door advising visitors of necessary actions.											
<b>All Healthcare Workers (HCWs)</b> on the ward are:		<ul style="list-style-type: none"> <li>Aware of the status of the ward and how Norovirus is transmitted.</li> <li>Aware of their duty to report when they have symptoms of gastrointestinal infection and not return or continue to work until they are symptom free for 48 hours.</li> </ul>									
<b>All patients</b> (and relatives) on the ward are aware of the Norovirus situation and have been given information leaflets on Norovirus and advised on the need for hand hygiene, and safe handling of personal laundry.											
<b>All patients</b> with symptoms of Norovirus have been assessed today for symptom severity and assessed for signs of possible dehydration (Stool and Fluid Balance charts).											
<b>Norovirus Outbreak Data Record</b> (Appendix 3). The outbreak data collection record has been updated – including any new cases, the symptoms patients are experiencing today and laboratory data. (Stool/vomit samples have been requested from all symptomatic patients).											
<b>Patient Placement Assessment:</b> A patient placement assessment and any advised/suggested moves have been made today.											
<b>Personal Protective Equipment (PPE)</b> – gloves, apron, surgical (mask/visor – if risk of facial contamination with aerosols). There are sufficient supplies of PPE in the ward and staff are using it appropriately.											
<b>Hand hygiene is being carried out with liquid soap and warm water</b> – this can be followed by alcohol based hand rub.											
<b>Hand hygiene:</b> Patients are encouraged and given assistance to perform hand hygiene before meals and after attending the toilet.											
<b>Environment:</b> The environment is visibly clean including curtains and there is increased cleaning which includes decontamination of frequently touched surfaces with detergent and 1000ppm av cl. (cleaning records are up-to-date).											
<b>Environment:</b> There are no exposed foods in the ward area – even if unexposed all fruit should be washed before eating.											
<b>Equipment:</b> Where possible single patient use equipment is used and communal patient equipment avoided. All re-usable equipment is decontaminated after use. There are sufficient other sundries on the wards to enable the control measures to be implemented.											
<b>Linen:</b> Whilst the ward remains closed, categorise all discarded linen as “infected”.											
<b>Spillages:</b> All faecal and vomit spillages are decontaminated by staff wearing PPE. The spillage is removed with paper towels, and then the area is decontaminated with an agent containing 1000 pp, av cl. All waste arising is discarded as healthcare waste. PPE is then removed and hands washed with liquid soap and warm water.											
<b>Advice and Guidance:</b> HCWs have access to and follow NHS Board guidance on:		<ul style="list-style-type: none"> <li>The decontamination of body fluid spills, equipment, soft furnishings.</li> <li>What to do if uniforms become contaminated.</li> </ul>									
<b>Today the ICT</b> has made an assessment of the outbreak and the continuing need for ward closure.											
<ul style="list-style-type: none"> <li><b>In preparation for re-opening</b> – empty beds have been cleaned but left unmade.</li> <li><b>In preparation for re-opening</b> – the curtains/bed screens in empty rooms have been taken down.</li> <li><b>In preparation for re-opening</b> – consider if pre-booking a terminal clean and pre-booking clean curtains/screens being hung is possible.</li> <li><b>Before re-opening:</b> a terminal clean has been performed following ICT recommendation and following the hospital procedure.</li> </ul>											

