



Think Twice!

This patient has a PVC in situ. All intravascular devices are associated with an increased risk of phlebitis, thrombosis, local infection and *S. aureus* bacteraemia. It is essential that you justify the need for your patient to have a PVC daily using the DRIFT mnemonic.

- **Diagnostics** : Does the patient need the cannula for a diagnostic procedure e.g. CT scan
- **Resuscitation**: Is the patient at risk of cardiac or respiratory arrest?
Intravenous: Does the patient require IV medication? Could these be switched to oral?
- **Fluids**: Does the patient require intravenous fluids? Could this be switched to oral fluids?
- **Transfusion**: Does the patient require a transfusion of blood products?

If your patient does not meet any of the above criteria please consider removing the PVC.



Think Twice!

This patient has a PVC in situ. All intravascular devices are associated with an increased risk of phlebitis, thrombosis, local infection and *S. aureus* bacteraemia. It is essential that you justify the need for your patient to have a PVC daily using the DRIFT mnemonic.

- **Diagnostics** : Does the patient need the cannula for a diagnostic procedure e.g. CT scan
- **Resuscitation**: Is the patient at risk of cardiac or respiratory arrest?
Intravenous: Does the patient require IV medication? Could these be switched to oral?
- **Fluids**: Does the patient require intravenous fluids? Could this be switched to oral fluids?
- **Transfusion**: Does the patient require a transfusion of blood products?

If your patient does not meet any of the above criteria please consider removing the PVC.