


TITAN CHLOR PLUS[®]


BLOOD SPILLS



Wear disposable gloves and apron. When using the solution, safety glasses should be worn if there is a risk of splashing.



To make solution, dissolve TITAN Chlor Plus tablets (according to dilution instructions below) in cold water. Leave lid off dilution bottle until tablets have dissolved. Mark date and time prepared on dilution bottle. Place disposable paper with solution over blood. **Leave for 2 minutes** and dispose as clinical waste.



Clean and disinfect the area larger than the spillage area with TITAN Chlor Plus solution.



Keep solution for no longer than 24 hours. Dispose of remaining solution as healthcare waste. Rinse with running water.



Dispose of gloves and apron in orange waste stream. Wash hands after removing gloves and apron.

DILUTION INSTRUCTIONS

DISINFECTION OF BLOOD SPILLS (10,000 PPM AVAILABLE CHLORINE)			CONTACT TIME
Disinfection of Blood Spills using TITAN CHLOR PLUS 10,000ppm Solution	TITAN Chlor Plus Tablets	1 tablet per 100ml cold water	e.g. 5 tablets per 500ml
	Leave lid off dilution bottle until tablets have dissolved		e.g. 10 tablets per 1,000ml (1L)
			2 minutes
			2 minutes

WARNINGS & PRECAUTIONS



AVOID contact with skin and eyes. **DO NOT** breathe dust/fume/gas/mist/vapors/spray.



DO NOT use directly on urine or vomit. Mixing this product with acid or ammonia releases chlorine gas.



AVOID prolonged contact with stainless steel or clothing.



ALWAYS dispose used materials as clinical waste. Wash hands thoroughly after handling.




ALWAYS keep container tightly closed. Store in a secure dry place in suitable labelled containers.



ALWAYS keep out of reach of children.



ALWAYS dispose of each solution after 24 hours. Refer to your SDS and hospital policy.



USE ONLY with adequate ventilation.