
TITAN SANITISING GRANULES®

WET BLOOD SPILLAGE







Wear disposable gloves and apron. When using the granules, safety glasses should be worn if there is a risk of splashing.




Dispense TITAN Sanitiser granules directly on to the wet blood spill and **leave for 2 minutes.**



Use disposable paper to remove granules and dispose of as healthcare waste.



Clean the area larger than the spillage area with general purpose detergent.



Keep solution for no longer than 24 hours. Clearly mark date and time on the bottle. dispose of remaining solution as healthcare waste.



Dispose of gloves and apron ias healthcare waste. Wash hands after removing gloves and apron.

DILUTION INSTRUCTIONS

DISINFECTION OF WET BLOOD SPILLAGES

CONTACT TIME

Disinfection of large wet blood spills using **TITAN Sanitiser Granules**

TITAN Sanitiser Granules

No dilution required

e.g. 5 tablets per 500ml

2 minutes

WARNINGS & PRECAUTIONS



AVOID contact with skin and eyes. DO NOT breathe dust/fume/gas/ mist/vapors/ spray.



DO NOT use directly on urine or vomit. Mixing this product with acid or ammonia releases chlorine gas.



AVOID prolonged contact with stainless steel or clothing.



ALWAYS dispose used materials as clinical waste. Wash hands thoroughly after handling.



ALWAYS keep container tightly closed. Store in a secure dry place in suitable labelled containers.



ALWAYS keep out of reach of children.



ALWAYS dispose of each solution after 24 hours. Refer to your

policy.

SDS and hospital



USE ONLY with adequate ventilation.



PLEASE REFER TO SDS AND HOSPITAL POLICY