

STANDARD OPERATING PROCEDURE (SOP) Twice Daily Clean of Isolation Rooms

Page	1 of 4
Effective from	Dec 2022
Review date	Dec 2024
Version	8

The most up-to-date version of this SOP can be viewed at the following web page: www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control

This SOP applies to all staff employed by NHS Greater Glasgow & Clyde and locum staff on fixed term contracts and volunteer staff.

SOP Objective

To minimise the risk of healthcare associated infection (HAI) from the environment.

KEY CHANGES FROM THE PREVIOUS VERSION OF THIS SOP

Document Control Summary

Approved by and date	Board Infection Control Committee 15 th December 2022	
Date of Publication	30 th January 2023	
Developed by	Infection Control Policy Sub-Group	
Related Documents	National Infection Prevention and Control Manual	
	NHSGGC Decontamination Guidance	
	NHSGGC Hand Hygiene Guidance	
	NHSGGC Cleaning of Near Patient Equipment SOP	
	NHS Scotland National Cleaning Services Specification	
Distribution/Availability	NHSGGC Infection Prevention and Control web page	
	www.nhsggc.scot/hospitals-services/services-a-to-	
	z/infection-prevention-and-control	
Lead Manager	Director Infection Prevention and Control	
Responsible Director	Executive Director of Nursing	



Page 2 of 4 Effective from Dec 2022 Review Dec 2024

STANDARD OPERATING PROCEDURE (SOP) TWICE DAILY CLEAN OF ISOLATION ROOMS

Review Dec 2024 date Version 8

The most up-to-date version of this SOP can be viewed at the following web page: www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control

Aim	To minimise the risk of healthcare associated infection (HAI) from the environment.	
Statement	Patients with certain alert organisms / conditions must be nursed in isolation to prevent the spread of the infection to other patients and staff. This is known as source isolation . In some situations if more than one patient is affected, during an outbreak of vomiting and diarrhoea for example, the restrictions may apply to a whole ward or part of a ward.	
Communication	The ward / area will notify the domestic staff when twice daily cleaning is required. An isolation notice will be displayed at the entrance to a single room/cohort area	
Requirements	The Domestic Supervisor / Team Leader will ensure domestic staff have the necessary equipment and that staff working in the area are fully trained on the correct procedures.	
	 Disposable or launderable mop (yellow) head Appropriately colour coded re-usable items (yellow), e.g. bucket and / or small bucket and bowl Dust control mop (disposable cover) Single-use cloths Single-use scourer Non-sterile single-use gloves Yellow single-use apron Clinical waste bag (orange) Chlorine based detergent (1000ppm) Chlorine based solution for sanitary ware Wet floor sign 	
	Where possible it is intended that individual cleaning equipment should be used per isolation room unless there are two or more in an individual bay, then the same isolation equipment may be used but water should be changed every 15 minutes. Personal Protective Equipment (PPE, i.e. gloves, aprons) must be changed between each room / bed bay	
	All re-usable equipment must be cleaned with chlorine based detergent.	
	If multiple isolation rooms are being cleaned, the solution must be	



Page 3 of 4 Effective from Dec 2022 Review date Dec 2024 Version 8

STANDARD OPERATING PROCEDURE (SOP) Twice Daily Clean of Isolation Rooms

The most up-to-date version of this SOP can be viewed at the following web page: www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control

Procedure	changed between rooms or after 15 minutes, whichever comes first. Cloths used in sanitary areas must be discarded immediately after use and a fresh cloth must be used for the general area. There should be a minimum of 4 hours between cleans. The responsibility for specific cleaning of equipment within a room may be			
Procedure	dependent on ward, department or specialty and therefore staff should check local policy as well as the SOP for Cleaning of Near Patient Equipment.			
	NB: please see SOP Cleaning of Near Patient Equipment			
	Prior to entering a room / area: Domestic Services Staff Report to the nurse in charge to ensure that it is convenient for cleaning to start and to receive any special instruction (additional PPE).			
	 Make up solution of chlorine based detergent (1000ppm) as per manufacturer's guidance Ensure you have the correct solution for cleaning sanitary ware Collect any equipment and materials required for cleaning inside the room / area as stated above. Carry out hand hygiene using alcohol hand gel or liquid soap and 			
	 water. Apply appropriate PPE, put on a disposable yellow plastic apron and disposable gloves (check with the nurse in charge if any other protective clothing is required). 			
	Procedure within isolation room: The door to the room must remain closed until the following procedure has been completed unless patient safety is compromised with the door closed (nursing staff will advise). PPE should be removed and disposed of as clinical waste and hand hygiene carried out prior to leaving the room / area. (NB FFP3 masks/reusable face visors should be removed prior to leaving the room / area and hands decontaminated thereafter)			
	 Explain to the patient what you are going to do in the room. Check room / area and if there is any visible contamination with 			



STANDARD OPERATING PROCEDURE (SOP) TWICE DAILY CLEAN OF ISOLATION ROOMS

Page	4 of 4	
Effective from	Dec 2022	
Review date	Dec 2024	
Version	8	

The most up-to-date version of this SOP can be viewed at the following web page: www.nhsggc.scot/hospitals-services/services-a-to-z/infection-prevention-and-control

Procedure (cont/ ...)

blood / body fluids inform nursing staff who will decontaminate the area with chlorine based detergent (10,000ppm) before you commencing cleaning.

- Gather items of debris and place in a clinical waste bag (orange).
- Clinical waste bags should be sealed in accordance with the NHSGGC Waste Policy.
- If the isolation room has been used for a patient with CDI, then chlorine based detergent must have a minimum contact time of 10 minutes for all surfaces and equipment NB: please see <u>SOP Cleaning</u> of Near Patient Equipment
- Replenish supplies (e.g. paper towels, soap) within the room.
- If there is a hand wash basin or en suite facilities ensure hand wash basin, shower, tiles, all fixtures and fittings, toilet seat and toilet bowl are thoroughly cleaned using a chlorine based detergent for use on sanitary ware (1000ppm). Always clean the toilet last and dispose of cloth immediately into orange clinical waste bag.
- A mop used to clean an isolation room should be used in 1 room and then either discarded as clinical waste or laundered after use.
- No buffing should take place in the room
- Replace waste bags.
- Once clean is complete, cleaning equipment should be removed as follows:

For Vermop:

The mop head should be bagged within the room for laundering, the mop handle wiped down with a solution of 1,000ppm and both removed from room

For mop and bucket system:

- The mop bucket should be emptied down the WC within the ensuite or taken to the DSR for emptying.
- The mop head bagged for laundering, the mop handle and bucket wiped with a solution of 1,000ppm and all removed from room.

Remove PPE and dispose of as clinical waste and undertake hand hygiene.